

CAPSUS INCISURATUS.

Capsus incisuratus, *Walk. Cat. Hem. Het. B. M.* vi. p. 121, n. 282 (1873).

Described from Ceylon.

CAPSUS ALBIPES.

Capsus albipes, *Motsch. Bull. Mosc.* xxxvi. (3) p. 82 (1863).

Described from Ceylon (Patannas).

*†CAPSUS RAVANA, sp. n. (Plate IV. fig. 10.)

Long. corp. 7 millim.

Testaceous, more or less mottled and speckled with reddish; head and pronotum with longitudinal reddish or brownish stripes; pronotum longitudinally striated, with the shoulder-angles rather prominent; scutellum punctured, brownish, with three pale longitudinal lines meeting at the tip; tegmina and wings hyaline; carina with the nervures yellowish towards the base and red towards the extremity, enclosing a large yellow space, of a long triangular form, on the costa at the extremity of the corium. Antennæ reddish, thickened towards the extremity of the second joint, and the terminal joints darker, as are likewise the tarsi.

Pundaloya, very common.

Somewhat approaches the genus *Lopus*.

*†CAPSUS RAMA, sp. n.

Long. corp. 8-10 millim.

Yellow, vertex with a slender black line between the eyes, and meeting behind them; pronotum blackish at the base, and with narrow black central and marginal lines, or with three short black lines at the base, the central line reddish, and the lateral lines reddish, black only at the base; pale part of the pronotum transversely striated, the hinder margin black, the lateral angles not produced. Scutellum transversely striated, more or less black towards the base and extremity, and divided by a deep groove. Wings hyaline, the corium with brown nervures; the costal nervure, and the opaque space at the extremity of the corium reddish, the latter yellowish in the centre. Antennæ reddish brown, darker or lighter, the second joint not distinctly thickened, joints 3 and 4 narrowly yellow at base. Legs yellowish, tarsi black, hind