

femora dotted with brown, hind tibiæ red. Under surface of body yellow, with a narrow red line on each side.

Pundaloya.

Allied to *Capsus lineifer*, Walk.

\*†CAPSUS ANTENNATUS, sp. n.

Long. corp. 7-8 millim.

Head and pronotum testaceous, finely punctured, pronotum frequently varied with brown in the middle or behind; antennæ with the last three joints more or less black at the extremities, and the second often black in the middle as well; scape sometimes likewise black; scutellum brown, tip usually testaceous; corium brown, set with short white bristles; costa broadly testaceous, intersected by a large reddish-brown spot just before the extremity; legs testaceous, more or less dotted with brown, and with two obsolete brown bands towards the tip of the hind femora; legs with longer bristles than the other Cinghalese *Capsi*. Pectus blackish in the middle; ventral surface of the abdomen variable in colour, reddish, testaceous, or blackish.

Pundaloya.

\*†CAPSUS LANKANUS, sp. n.

Long. corp. cum tegm. 4 millim.

Head and thorax black, antennæ with the base of the first and third joints white; eyes white, clavus black; corium chestnut-brown, with a white transverse band near the base, and another preceding the tip, which is blackish; membrane and wings subhyaline; abdomen beneath with a broad white transverse band at the base.

The type is from Nitagala.

DERÆOCORIS RUBROVULNERATUS.

*Deræocoris rubrovulneratus*, *Motsch. Bull. Mosc.* xxxvi. (3) p. 83 (1863).

Described from Ceylon (Nura Ellia).

DERÆOCORIS VIRIDANUS.

*Deræocoris viridanus*, *Motsch. Bull. Mosc.* xxxvi. (3) p. 83 (1863).

Described from Ceylon (Nura Ellia).

DERÆOCORIS PICEONIGER.

*Deræocoris piceoniger*, *Motsch. Bull. Mosc.* xxxvi. (3) p. 84 (1863).

Described from Ceylon (Colombo).