

- Genus 92.—**Plexaris** Kirkaldy, 1902, Entomologist, xxxv, 282. Type **satur-nides** Kirkaldy.
- Genus 93.—**Platycapsus** Reuter, 1904, Ofv. Finsk. Vet. Förh., xlvii, No. 4, 11 (sep. ?). Type **acaciae** Reuter.
- Genus 94.—**Deracocoris** Kirschbaum, 1855,† J. b. Ver. Nat. Nassau, x, 208 (sep. 31 !) (type fig. in Wolff, Icones, Cimic., i, pl. 4. f. 35) = || *Capsus* of many authors (nec Fabricius typ.) = *Macrocapsus* Reut., 1879, Ofv. Finsk. Vet. Förh., xxi, 55 (type fig. in Stål, Stett. Ent. Zeit., xix, pl. 1, f. 1). Type *brachialis* (Stål), Reuter.† Type **medius** Kirschb. (= *tricolor* Fabr.), Dist.
- Genus 95.—**Chilocrates** Horváth, 1889, Termész. füz., xii, 39. Type **lenzii** Horváth.
- Genus 96.—**Shana** Kirkaldy, 1902, Entomologist, xxxv, 316. Type **ravana** Kirkaldy.
- Genus 97.—**Saundersiella** Reuter, 1890, Rev. Ent., ix, 262 = || *Saundersia* Reuter, 1876, Pet. nouv. ent., ii, 32 (type fig. in Reuter, Hem. Gymn. Eur., v, pl. 1, f. 6†). Type **moerens** Reuter.
- Genus 98.—**Romna** Kirkaldy, n. n., = || *Morna* F. B. White, 1878, Ent. Mo. Mag., xv, 130. Type **capsoides** White.
- Genus 99.—**Coccobaphes** Uhler, 1878, P. Boston N. H. Soc., xix, 461 = † *Caccobaphes* Reuter, 1879, Zool. Jahrb., 507. Type **sanguinarius** Uhler.
- Genus 100.—**Kalanía** Kirkaldy, 1904, Entom., xxxvii, 280 = || *Baracus* Kirkaldy, 1902, Fauna Hawaii., iii, 143. pl. 4, f. 21.† Type **hawaiien-sis** Kirkaldy.
- Genus 101.—**Stethoconus** Flor, 1861, Arch. Naturk. Kurlands (2), iv (sep. ?), 615 = *Acropelta* Mella, 1869, Bull. Soc. E. Ital., i, 292, pl. 4.† Type *pyri* Mella (= *cyrtopeltis* Flor). Type **cyrtopeltis** Flor.¹⁸
- Genus 102.—**Bothynotus** Fieber, 1864, Wien. Ent. Mon., viii, 76 = *Trichy-menus* Reuter, 1875, Not. Faun. Fenn., xiv, 7. Type *pilosus* (Boh.), Reuter. Type fig. in Reuter, Hem. Gymn. Eur., v, pl. 4. f. 1. Type **minki** Fieber (= *pilosus* Boheman).
- Genus 103.—**Alloetomus** Fieber, 1858, Wien. Ent. Mon., ii, 303 (type fig. in H. S., Wanz. Ins., iii, f. 284, as *marginepunctatus*). Type **gothicus** (Fall.), Fieber.
- Genus 104.—**Euarmosus** Reuter, 1875, O. V. A. F., xxii, 76. Type **sayi** Reuter.
- Genus 105.—**Eubatas** Distant, 1884, B. C. A. Het., i, 277. Type **chiriqui-nus** Distant, pl. 27, f. 8.†
- Genus 106.—**Neocapsus** Distant, l. c. Type **mexicanus** Distant, pl. 22, f. 19.†
- Genus 107.—**Horcias** Distant, l. c. Type **variegatus** Dist., pl. 24, f. 13.*†
- Genus 108.—**Calocorisca** Distant, op. c., 280. Type **villosa** Dist., pl. 24, f. 17.*†

¹⁸ Reuter, Atkinson and in fact all authors persist in calling this *mamillosus* (Flor), but I can see no reason for the rejection of the earlier *cyrtopeltis* Flor. They have also overlooked the fact that Flor, not Fieber, is the author of the genus.