

margins of hind area divergent; hind margins a little sinuate. Cuneus elongate. Membrane with two cells, the inner almost oblong, obliquely truncate apically.

37. *S. CHAMBERSI*, sp.nov. (Pl.xliiii., fig.8).

Shining orange-red; legs, etc., paler. Eyes blackish. Antennæ dark reddish-fuscous. Membrane hyaline, apically smoky, veins scarcely red. Wing-veins pale fuscous. Abdomen dark blood-red. Hind tibiæ with three pale fuscous rings. Head, pronotum, and scutellum not punctured. Antennæ 25, 100, 25, 23, the first segment half as long as the width of the pronotum apically. Hindmargin of pronotum three-fifths longer than the apical margin. Length 6 mill.

*Hab.*—Q.: Kuranda (Aug.; Perkins); one female in poor condition.

I have pleasure in naming this after Mr. W. E. Chambers, who has adorned my recent memoirs with many fine drawings.

PARACALOCORIS Distant.

38. *P. AUSTRINUS*, sp.nov.

Ochreous-brown, with pale golden-yellow pubescence. Head with a black impression just basal of the antennal insertions. First segment of the antennæ dark reddish-fuscous, second ochreous-fuscous with the apical half blackish, third dark fuscous with the base pale ochreous. Labium pale ochreous-fuscous, third segment and apex of the fourth blackish. Collar narrowly margined posteriorly with black. Pronotum with two pale smoky-fuscous broad bands down the middle of the hind area. The transversely impressed line on the scutellum blackish, anterolateral angles broadly pale smoky-fuscous. Corium mostly suffused with pale smoky-fuscous; cuneus subsanguineous, apical margin smoky. Commissural margin narrowly blackish. Membrane hyaline, apical margin irregularly dark smoky; veins blackish, apical angle of the inner cell suffused. Mesopleuræ each with a large blackish spot; underside generally more or less