

twice as wide as anterior margin; punctured and covered with long tufts of hair. Scutellum with two or three tufts. Labium extending to base of middle coxæ. Tegmina explanate laterally.

88. *N. KNOWLESI*, sp.nov.

Head, sterna, etc., testaceous. Antennæ dark fuscous; basal half and apex of first; base, a middle ring, and apex of second; apices of third and fourth; whitish-testaceous. Pronotum pale purplish-brown, two subanterior spots, collar, and posterior margin (very narrowly) pale. Scutellum, clavus and interior third of corium, dark purplish-brown; posterior angle of scutellum greenish-yellow. Rest of corium dark fuliginous, spotted with hyaline. Sternites yellowish-green. Legs multiannulate with fuscous. Length 5 mill.

Lautoka (Knowles, No.349), on *Artocarpus incisa* and Tobacco-leaves.

I hope to amplify this diagnosis later and figure the interesting Mirid.

Fam. CICADIDÆ.

89. *CYCLOCHILA AUSTRALASIE* (Donovan), Goding & Froggatt, 1904, Proc. Linn. Soc. N. S. Wales, xxix. 569.

Recorded by me (1907, Bull. Ent. Haw. Sugar Plant. Assoc. iii. 17), but Mr. Froggatt tells me that he thinks that the individuals were accidental visitors, or introduced into Fiji, as this species has a very limited range in Australia.

90. *MACROTRISTRIA ANGULARIS* (Germar).

*Cicada angularis* Goding & Froggatt, 1904, *op. cit.* 580.

The same remarks apply to this as to the former.

91. *CICADA KURUDUADUA* Distant, Trans. Ent. Soc. London, 1881, 645; Waterhouse, 1882, Aid Ident. Ins. i. Pl. C.

"Fiji."

92. *C. KNOWLESI* Distant, 1907, Ann. Mag. Nat. Hist.(7) xx. 413.

"Fiji."

93. *COSMOPSALTRIA VITIENSIS* (Distant).

*Sawda*(?) *vitiensis* Distant, 1906, Entom. xxxix. 12.

"Fiji."