

Pronotum: Length .55 mm., width at base 1.11 mm.; black, scarcely shining. Scutellum black. Sternum, pleura, and ostiolar peritreme, dull black.

Hemelytra: Embolar margins very slightly arcuate; fuscous black, base of cuneus scarcely paler than corium; yellowish pubescent, hairs dusky on cuneus and to some extent on embolium. Membrane uniformly fuscous, veins pale, spot bordering apex of cuneus clear.

Legs: Dark fuscous, apices of femora slightly yellowish; small opaque black spots visible on anterior and posterior faces of femora; tibiae fuscous but black spots at bases of spines showing through the infuscation. Venter black, covered with an opaque bloom.

*Female*: Length 3.6 mm., width 1.55 mm.; more ovate and robust than the male but very similar in coloration.

Food plant: *Larix laricina*.

Holotype: Male, 27 June, 1920, Ithaca, N. Y. (H. H. Knight); author's collection. Allotype: same data as type. Paratypes: Males and females (64), taken with the types on *Larix laricina*. CONNECTICUT—Male, 7 July, 1920, New Haven (B. H. Walden). MAINE—Female, 12 July, Princeton (C. W. Johnson). Male, 5 July, 1911, Orono. MINNESOTA—Males and females (36), 12 Aug., 1922, Beaver Dam, Cook County (H. H. Knight). NEW YORK—Males and females (18), 26 July, 1916, Ithaca (H. H. Knight). ALBERTA—Males (2) and females (3), 4 Aug., 1921, Nordegg (J. McDunnough).

*P. repetitus* Knight, new species.

Smaller than *laricicola* but the dorsal aspect very similar; legs yellowish, femora with small, rather inconspicuous fuscous dots arranged in series on anterior face.

*Male*: Length 3 mm., width 1.24 mm. Head: Width .64 mm., vertex .33 mm.; black, shining, vertex scarcely paler. Rostrum, length .55 mm., yellowish, basal and apical segments blackish.

Antennae: Segment i, length .22 mm., black; ii, .75 mm., black; iii, .55 mm., fuscous; iv, .39 mm., pale fuscous.

Pronotum: Length .53 mm., width at base .97 mm.; black. Scutellum, sternum, pleura, and ostiolar peritreme black, moderately shining.

Hemelytra: Embolar margin only very slightly arcuate; black, moderately shining, clothed with fine yellowish pubescence; cuneus uniformly colored like the corium, not at all paler at fracture. Membrane and veins uniformly pale fuscous, scarcely paler bordering tip of cuneus.

Legs: Yellowish, coxae fuscous on basal half; femora spotted with fuscous, arranged in one or two rows on anterior face; tibiae with rather prominent black spots at bases of spines. Venter black, shining, minutely yellowish pubescent.

*Female*: Length 2.9 mm., width 1.16 mm.; very similar to the male in form and coloration.