

Antennae: Segment i, length .30 mm., yellow, fuscous at base; ii, .86 mm., yellow, fuscous at base and dusky on apex; iii, .47 mm., yellowish and tinged with dusky; iv, .33 mm., pale fuscous.

Pronotum: Length .55 mm., width at base 1.11 mm.; blackish, moderately shining; central area of disk and the basal angles pale to yellowish; clothed with fine yellowish pubescence. Scutellum pale yellowish, median line rather broadly, and the mesoscutum, black. Sternum, pleura, and ostiolar peritreme, black.

Hemelytra: Embolar margins slightly arcuate; blackish, embolium, cuneus, and basal half of corium, pale to yellow, sometimes distinctly reddish; clothed with fine yellow pubescence, longest on base of clavus. Membrane fuscous, paler bordering apex of cuneus, veins pale only at apex of areoles.

Legs: pale to yellowish, basal half of hind coxae and tips of tarsi fuscous. Venter blackish, moderately shining.

*Female*: Length 3.4 mm., width 1.39 mm.; slightly more robust than the male but very similar in coloration.

Food plant: *Quercus alba*.

Holotype: Male, 11 June, 1919, University Farm, St. Paul, Minn. (H. H. Knight); Minn. Univ. Coll. Allotype: taken with the type. Paratypes: Males and females (12), taken with the types on *Quercus alba*. Males and females (8), 10 June, 1920, type locality (H. H. Knight). NEW YORK—Male, 25 June, 1915, Batavia; females (2), 23 June, 1916, Conesus Lake (H. H. Knight).

*M. maculipennis* var. *fuscicornis* Knight, new variety.

Similar to the typical *maculipennis* but distinguished by the black antennal segments, segment i narrowly yellowish at apex.

Holotype: Male, 29 June, 1905, Monmouth, Maine (C. W. Johnson); author's collection.

*M. nigricornis* Knight, new species.

Black, strongly shining, minutely pubescent; antennae yellow, segment ii and base of i, black; legs pale.

*Male*: Length 4 mm., width 1.44 mm. Head: width .66 mm., vertex .33 mm.; black, vertex slightly pale. Rostrum, length 1.22 mm., yellowish, basal segment and apex blackish.

Antennae: Segment i, length .25 mm., yellowish, black at base; ii, .94 mm., deep black, narrowly pale at base; iii, .69 mm., pale yellow; iv, .36 mm., pale yellowish.

Pronotum: Length .61 mm., width at base 1.11 mm.; black, minutely pale or dusky pubescent. Scutellum, sternum, pleura, and ostiolar peritreme, black.

Hemelytra: Embolar margins nearly straight; uniformly black, strongly shining; dusky or black pubescent. Membrane and veins uniformly fuscous, not or scarcely paler bordering tip of cuneus.

Legs: Pale to yellowish; bases of coxae and tips of tarsi black-