

## Teleorhinus Uhler.

*T. tephrosicola* Knight, new species.

*Male*: Length 7.3 mm., width 2.2 mm.; differs from *cyaneus* Uhler in that the combined length of antennal segments iii and iv is greater than the length of segment ii; clavate portion of segment ii only twice as thick as segment i, while in *cyaneus* the apical half of segment ii is three times the thickness of segment i.

Antennae: Segment i, length .36 mm., black; ii, 2.05 mm., apical two-fifths clavate, .17 mm. thick, yellowish, the clavate portion black; iii, 1.39 mm., slender, fuscous, slightly paler at base; iv, .97 mm., fuscous.

Black, shining, minutely pubescent, very similar to *cyaneus* but differs in form of anterior angles of pronotum; legs pale reddish yellow, coxae whitish but blackish at base; rostrum brownish but darker at base and apex.

*Female*: Length 7.6 mm., width 2.5 mm.; not differing materially in form or color from the male.

Holotype: Male, 11 July, 1916, Yaphank, Long Island, N. Y. (Wm. T. Davis); author's collection. Allotype: taken with the type. Paratype: Female, 17 June, 1906, Lakehurst, N. J. (Wm. T. Davis). Mr. Davis took the type specimens on flowers of *Tephrosia* sp.

## Subfamily DICYPHINAE.

*Key to Genera.*

1. Pseudarolia prominent (fig. 49: 18-20) ..... 2  
Pseudarolia absent (fig. 49: 21); hemelytra hyaline, glassy, ovate, with a sharply defined inverted Y-shaped red or fuscous mark (p. 478) *Hyaliodes*
2. Calli distinctly impressed at basal margin, and thus the pronotal disk rather distinctly transversely sulcate ..... (p. 476) *Dicyphus*  
Calli not or scarcely impressed at basal margin, pronotal disk not transversely sulcate ..... 3
3. Eyes small, head elongated behind eyes for a space equal to lateral width of an eye ..... (p. 478) *Macrolophus*  
Eyes large, head space behind eyes not equal to half the lateral width of an eye ..... [*Cyrtopeltis*]

*Dicyphus* Uhler.*Key to Species.*

1. Ostiolar peritreme strongly protruding, ostiole large and clearly exposed ..... 2  
Ostiolar peritreme scarcely raised from the pleural wall, ostiole inconspicuous or quite concealed; head and pronotum black, collar and ray between calli pale, hemelytra chiefly pale ..... *agilis*
2. Eyes removed from pronotal collar for a space equal to dorsal width of an eye; hemelytra conspicuously colored with red .... *famelicus*  
Eyes removed from pronotal collar for a space equal to less than dorsal width of an eye; hemelytra darkened with fuscous, rarely tinged with reddish ..... *discrepans* n. sp.