

Female: Length 4.6 mm., width 1.84 mm.; very similar to the male in form and color.

Holotype: Male, 14 July, 1916, Batavia, N. Y. (H. H. Knight); author's collection. Allotype: same data as the type. Paratypes: Males and females (30), taken with the types on hickory. GEORGIA—Male and female, July, 1910, Rabun County (Wm. T. Davis). MARYLAND—Male, 21 June, 1914, Plum Point (W. L. McAtee). MINNESOTA—Female, 20 July, 1920, Gray Cloud Island (H. H. Knight). NORTH CAROLINA—Female, 14 June, 1915, Southern Pines (A. H. Manee). Males and females (8), July, 1912, Black Mountains (Beutenmuller). NEW JERSEY—Female, 2 July, Jamesburg (Wm. T. Davis). NEW YORK—Males and females (7), 17 July, males and females (12), 31 July, female, 6 Aug., 1916, females (2), 13 Aug., 1915, Batavia (H. H. Knight), collected on hickory. Males and females (5), 16 July, 1916, Conesus Lake; males and females (6), 23 July, 1916, Ithaca (H. H. Knight). Female, 7 Aug., 1915, White Plains (J. R. T. Bueno). Males (4) females (3), 30 June, 1922, Kings Bluff, Winona Co., Minn. (H. H. Knight).

Collected most frequently on hickory (*Carya* sp.). Nymphs have been taken on the trunks and larger limbs of trees, indicating that the species may be largely predaceous.

C. vicinus Knight, new species.

Fulvous to reddish, scutellum and apical half of hemelytra dark brownish to blackish, legs and antennae yellowish; dorsum clothed with rather inconspicuous simple pubescence, scutellum and basal half of hemelytra thickly clothed with closely appressed, scale-like silvery pubescence.

Male: Length 3.4 mm., width 1.4 mm. Head: Width .78 mm., vertex .31 mm.; yellowish to reddish, front coarsely alutaceous, vertex and front beset with a few erect yellowish hairs. Rostrum, length 1.43 mm., reaching upon hind coxae, reddish, yellowish apically.

Antennae: Segment i, length .31 mm., yellowish, frequently a red mark on ventral surface; ii, 1.23 mm., thickness at apex .10 mm., yellowish, more brownish at apex; iii, .66 mm., thickness .08 mm., reddish brown, paler at base; iv, .48 mm., reddish brown.

Pronotum: Length .71 mm., width at base 1.23 mm.; lateral margins of disk nearly straight; fulvous to reddish, moderately shining; calli outlined but inconspicuous, alutaceous; sparsely clothed with fine yellowish pubescence. Scutellum dark reddish brown, transversely rugulose, thickly clothed with closely appressed, silvery scale-like pubescence. Sternum and pleura reddish to brownish, meta-episterna pruinose; ostiolar peritreme reddish.

Hemelytra: Embolar margins slightly sinuate; yellowish to fulvous, apical half dark brownish to blackish; sparsely clothed with simple yellowish pubescence, clavus and basal half of corium covered with closely appressed, silvery scale-like pubescence. Membrane pale, apical half fuscous.

Legs: Uniformly yellowish, scarcely darker at base of coxae. Venter dark reddish brown to black, shining, genital segment