

ostiolar peritreme yellowish, dorsal lobe strongly protruding, fuscous just above.

Hemelytra: Embolar margins nearly straight; yellowish translucent, apical half of corium and the cuneus darkened with fuscous; finely pubescent and interspersed with erect, yellowish pilose hairs, pubescent hairs apparently arising from fine punctures. Membrane pale, a large somewhat rounded fuscous cloud occupying central area between areoles and including the apical half.

Legs: Pale yellowish brown, front coxae reddish to fuscous. Venter reddish brown to blackish, darker distally, shining, clothed with yellowish pubescence; genital claspers distinctive.

Female: Length 5 mm., width 1.43 mm.; slightly more robust than the male but very similar in coloration, embolar margins more arcuate distally.

Holotype: Male, 15 July, 1915, Parry Sound, Ontario, Canada (H. S. Parish); author's collection. Allotype: same data as the type. Paratypes: Male, taken with types. MICHIGAN—Male, 22 July, 1916, Pentwater (E. Liljeblad). NEW HAMPSHIRE—Female, Franconia (Mrs. A. T. Slosson). QUEBEC—Male, 11 July, 1920, Cascades (H. G. Crawford). SOUTH DAKOTA—Male, 16 July, 1920, Brookings (H. C. Severin).

C. setosus Reuter.

Acta Soc. Sci. Fenn., xxxvi, No. 2, 70, 1909.

Female: Length 2.5 mm., width 1.3 mm.; dark brownish with yellowish; distinguished by the small size and oval shape; hemelytra shorter than the abdomen, membrane absent, tips broadly rounded, finely and closely punctured, clothed with short, closely appressed, sericeous pubescence and interspersed with long erect setose hairs.

Male: Length 3 mm., width 1.3 mm.; wings fully developed, suggestive of *pumilus* but smaller; distinguished by the small size and erect setose hairs on dorsum.

New Jersey, Maryland, North Carolina.

Pamillia Uhler.

P. davisii Knight, new species.

Distinguished from *behrensi* by the glabrous and strongly shining area on apical one-third of corium, also by the dark reddish brown color.

Female: Length 3.8 mm., width 1.5 mm. Head: Width .86 mm., vertex .44 mm.; front more nearly flat and more sharply declivitous than in *behrensi*, in form very similar to certain species of *Pilophorus*; dark reddish brown, nearly black on vertex, front and vertex alutaceous, lower half of face more strongly shining; finely pubescent, a few longer hairs on front. Rostrum, length 1.36 mm., scarcely attaining hind margins of intermediate coxae, dark reddish brown.