

P. furvus Knight, new species.

Female: Length 4.1 mm., width 1.61 mm. Head: Width 1.12 mm., vertex .61 mm., basal margin to tip of tylus .89 mm.; dark reddish brown to piceous, smooth, shining; gula beset with erect, bristly pubescent hairs. Rostrum, length 1.4 mm., scarcely attaining hind margin of sternum, rufo-piceous.

Antennae: Segment i, length .25 mm.; ii, 1.03 mm., thickness (.08 mm.) equal to segment i, more slender on basal one-fourth, minutely pubescent, dark reddish brown, slightly paler at base; iii, .41 mm., slender, pale, apical half dark brown; iv (missing).

Pronotum: Length .83 mm., width at base 1.33 mm., anterior angles .77 mm.; lateral margins sulcate, flaring sharply to basal angles, a distinct bristle just before each anterior angle; uniformly dark brownish black; xyphus convex, lateral margins scarcely carinated, reddish brown. Scutellum strongly convex, almost taking the form of a truncate pyramid but rounded above, apex flattened, basal angles and apex with a tuft of silvery scale-like pubescence; mesonotum broadly exposed, flattened above but abruptly declivitous at each side. Sternum brownish black, polished; epimera with posterior margin covered by silvery scale-like pubescence; ostiolar peritreme pale to dusky, polished just above, a tuft of scale-like pubescence at posterior margin of metasternum just beneath that on embolium.

Hemelytra: Embolar margins sulcate-sinuate, greatest width (1.61 mm.) just before cuneal fracture; from anterior to posterior silvery line, opaque, dark brownish black; behind posterior silvery line, brownish black to piceous, polished, apex of clavus and all of the cuneus included; posterior silvery line sinuate, perhaps interrupted (mutilated?), nearly or quite connecting with that of clavus which is directed slightly distad; basal silvery line similar to that of *amoenus*; very finely pubescent, longer on apical margin bordering membrane. Cuneus sharply deflected, piceous and polished similar to apex of corium, a silvery scale-like spot near inner basal angle. Membrane pale fuscous, a large opaque blackish cloud involving apical half of areoles and middle of membrane.

Legs: Dark brownish black, tibiae more yellowish brown, trochanters except on front legs, two basal segments of tarsi, and apical half of hind coxae, pale.

Venter: Dark brown to piceous, polished, finely pubescent, prominent bristly hairs bordering ovipositor.

Holotype: Female, 30 June, Lakehurst, N. J. (Wm. T. Davis); author's collection.

Pilophorus depictus Knight, new species.

Closely related to *furvus* but smaller, length of antennal segment ii scarcely equaling width of head; hemelytra, anterior to posterior silvery line, cinnamon fulvous as in *amoenus*.