

Pronotum: Length .94 mm., width at base 1.53 mm., anterior angles .83 mm.; anterior half of disk sparsely clothed with silvery, deciduous, pubescent hairs similar to those on front of head. Scutellum with apical half and slender lateral margins flat, abruptly convex on basal half but flattened basally, flattened apical half more or less covered with silvery, scale-like pubescence.

Hemelytra: Dark fusco-brownish, opaque anterior to posterior silvery line, basal half with a silvery sheen apparent in certain lights; beset with erect, short black bristles; posterior silvery line nearly straight, slender, behind this distinctly polished, apex of clavus included. Membrane uniformly darkened with fuscous, an opaque black cloud bordering apex of larger areole.

Legs: Uniformly brownish black, a pale spot on anterior aspect of front coxae near base; hind tibiae strongly flattened and distinctly curved. Venter with a patch of silvery, scale-like pubescence laterally on third segment.

*Female*: Length 5.3 mm., width 1.8 mm.; very similar to the male but antennal segment iii perceptibly thicker (.10 mm. thick), nearly equal to thickness of segment i.

Breeds on *Pinus resinosa*.

Holotype: Male, 11 July, 1920, Taghanic, N. Y. (H. H. Knight); author's collection. Allotype: same data as the type. Paratypes: Males and females (16), taken with the types on *Pinus resinosa*. MASSACHUSETTS—Female, 15 July-6 Aug., 1918, Woods Hole (Chris E. Olsen).

*P. uhleri* Knight, new species.

Closely related to the preceding species but differs in color and thickness of antennal segment iii, in having patches of silvery scale-like pubescence on sides of fourth and fifth abdominal segments, and by the fact that the hind tibiae are not so distinctly compressed.

*Male*: Length 5 mm. Head: Width 1.14 mm., vertex .53 mm., from tip of tylus to vertex .94 mm.; front without silvery scale-like pubescence. Rostrum, length 1.8 mm., nearly attaining hind margins of middle coxae.

Antennae: Segment i, length .38 mm., thickness .10 mm.; ii, 1.86 mm., gradually thickened apically (.14 mm. thick); iii, .66 mm., thickness .06 mm., pale, apical half infuscated, sometimes the whole segment tinged with pink; iv, .61 mm., pale, apex dusky.

Pronotum: Length .83 mm., width at base 1.44 mm.; uniformly black, beset with black pubescence but devoid of silvery scale-like hairs. Scutellum nearly as in *vanduzeei* but with sides and base thickly covered with silvery, scale-like pubescence.

Hemelytra: Very similar to *vanduzeei* but the posterior silvery line distinctly broader.

Legs: Hind tibiae slightly compressed but not distinctly flattened as in *vanduzeei*.

Venter: With a distinctive patch of silvery, scale-like pubescence on the sides, extending obliquely across segments 3-6 inclusive.