

apex; membrane fuscous, veins and central area paler, a small clear spot bordering apex of cuneus.

Breeds on *Galium boreale* in Minnesota.

Colorado, Minnesota, New York, Canada, Maine.

The typical *unifasciatus* is more broadly pale than variety *lateralis*, the embolium and corium pale except for a small fuscous patch on apical area of corium. Specimens of the typical form from North America have not been examined by the writer.

Polymerus punctipes Knight, new species.

Legs, embolium, and outer margins of cuneus, fulvous; hind femora with a small group of fuscous points on anterior face at middle of apical half.

Male: Length 4.6 mm., width 2 mm. Head: Width .97 mm., vertex .47 mm.; black, a yellow spot each side of vertex bordering eye; yellowish pubescent. Rostrum, length 1.23 mm., almost attaining hind margin of sternum, yellowish, apex blackish.

Antennae: Segment i, length .51 mm., black, narrowly yellowish at base; ii, 1.85 mm., nearly cylindrical, slightly thicker on middle, nearly equal in thickness to segment i, black, clothed with black and pale pubescence intermixed; iii, .66 mm., yellowish and tinged with dusky; iv, .80 mm., dusky.

Pronotum: Length 1 mm., width at base 1.7 mm.; black, slightly shining, disk transversely rugulose; clothed with pale and yellowish sericeous pubescence. Scutellum black, transversely rugulose, clothed with yellowish pubescence, more sericeous on basal angles. Sternum and pleura black; ostiolar peritreme yellow, becoming dusky on anterior lobe.

Hemelytra: Embolar margins slightly arcuate on apical half; black, embolium and outer margin of cuneus yellowish or fulvous; surface scabriculous, slightly shining, clothed with golden sericeous pubescence and intermixed with darker pubescent hairs. Membrane rather uniformly fusco-brownish, a small nearly clear spot bordering apex of cuneus, veins yellowish.

Legs: Fulvous, spot on knee and apical tarsal segment blackish; hind femora with a group of from two to five fuscous points on anterior face at middle of apical half, a prominent hair arising from each of the two lower points; pubescence pale to yellowish, but blackish on apical half of femora. Venter black, clothed with pale to yellowish pubescence.

Female: Length 5.2 mm., width 2.4 mm.; embolar margins strongly arcuate on apical half; pubescence and color similar to that of the male. Rostrum, length 1.31 mm., nearly attaining hind margin of sternum.

Holotype: Male, 4 July, 1915, Four Mile, N. Y. (H. H. Knight); author's collection. Allotype: same data as the type. Paratypes: DISTRICT OF COLUMBIA—Female, 12 June, 1906, Brightwood (O. Heidemann). FLORIDA—Female, "Jacksonville." MAINE—Male, 22 July, 1909, Machias (C. W.