In my key to the species of *Plagiognathus* in the "Hemiptera of Connecticut," *flavescens* runs to either *rosicola* or *blatchleyi*; but distinguished from *rosicola* by the much paler color, and from *blatchleyi* by the larger, more elongate form, and narrower vertex; width of vertex in the male is not equal to length of antennal segment I, or about equal in the female.

## Atractotomus acaciae n. sp.

Distinguished from albidicoxis and hesperius by the shorter antennal segment II which does not equal width of head, also

by the pale rostrum.

¿. Length 2.3 mm., width 1.14 mm. Head: width .66 mm., vertex .31 mm. Rostrum, length .84 mm., attaining middle of hind coxae, pale, last two segments black. Antennae: segment I, length .14 mm., thickness .085 mm.; II, .56 mm., thickness .128 mm., fusiform, clothed with prominent black pubescence; III, .199 mm., slender; IV, .185 mm.; black, last two segments pale. Pronotum: length .48 mm., width .98 mm. Black, tip of scutellum, basal angles of corium, and sometimes extending along claval suture, pale; thickly clothed above and below with white, closely appressed, scale-like pubescence, intermixed with fine dusky hairs, with two or three prominent black hairs at anterior angles of pronotum.

Q. Length 2.6 mm., width 1.3 mm. Head: width .67 mm., vertex .33 mm. Antennae: segment I, length .14 mm.; II, .51 mm., thickness .13 mm.; III, .199 mm.; IV, .185 mm. Very similar to the male in form and coloration; pale areas frequently more extensive, femora more or less pale and showing spots of blackish, tibiae paler with black spots at

base of spines.

Holotype: § June 7, 1924, Tucson, Arizona (A. A. Nichol); author's collection. Allotype: same data as the type. Paratypes:  $8 \$ \$ \$, taken with the types. 2 9 June 18, 1915, Lewis Spings, 9 June 18, Fort Huachuca, Arizona (Harold Morrison).

## Psallus suaedae n. sp.

Distinguished from other described species of *Psallus* by the fusco-reddish femora and dotted tibiae, combined with the pale greenish dorsum and venter.

8. Length 2.6 mm., width 1 mm. Head: width .71 mm., vertex .34 mm. Rostrum, length .77 mm., reaching upon middle of hind coxae. Antennae: segment I, length .17 mm.; II, .77 mm.; III, .51 mm.; IV, .34 mm.; with fine pale to