

♂. Length 3.6 mm., width 1.36 mm. *Head*: width .77 mm., vertex .19 mm. Rostrum, length 1.14 mm., reaching to middle of intermediate coxae. *Antennae*: segment I, length .23 mm.; II, .91 mm., cylindrical, equal in thickness to segment I; III, .57 mm.; IV, .47 mm.; pale yellowish, segment IV and apical half of III brown. *Pronotum*: length .60 mm., width at base 1.20 mm.; disk very minutely punctate, Dorsum clothed with very fine closely appressed sericeous, yellowish pubescence, intermixed with sparsely set, longer golden yellow pubescence, longest hairs on scutellum and clavus. Color castaneous brown, scutellum paler; hemelytra pale yellowish, apical half of corium and embolium, and the cuneus, fusco-brownish, red showing in disk of cuneus. Membrane pale yellowish, apical half and the anal area fuscous. Genital claspers very small, left clasper curving straight back to an acuminate point, right clasper very small, twice as long as broad, notched at apex, lower point longer.

♀. Length 3.6 mm., width 1.4 mm. *Head*: width .80 mm., vertex .34 mm. *Antennae*: segment I, length .28 mm.; II, 1.03 mm.; III, .66 mm.; IV, .54 mm. *Pronotum*: length .61 mm., width at base 1.23 mm. Very similar to the male in form, vestiture, and color.

Holotypes: ♂, July 27, 1917, Mt. Lemon, Santa Catalina Mts., Arizona, alt. 9,000 ft. (H. H. Knight); author's collection. *Allotypes* ♀, Aug. 3, 1915, Huachuca Mts., Arizona (H. G. Barber). *Paratypes*: ♂, taken with the type; Cornell University collection.

Ceratocapsus clavicornis n. sp.

Allied to *fusiformis* Van D., but distinguished by the differently formed genital claspers, and in color and type of pubescence; distinguished by the broad flattened blade on terminal portion of left genital clasper.

♂. Length 3.3 mm., width 1.3 mm. Rostrum, length 1.2 mm., reaching to middle of hind coxae. *Head*: width .74 mm., vertex .33 mm. *Antennae*: segment I, length .23 mm., thickness .086 mm.; II, .74 mm., slender at base and gradually thickened to apex (.09 mm. thick); III, .39 mm., thickness .10 mm., base more slender; IV, .41 mm., thickness .11 mm., dark brown, segment I and base of II yellowish brown, clothed with fine pale pubescence. *Pronotum*: length .71 mm., width at base 1.21 mm. Dorsal surface finely punctate as in *pumillus*, clothed with sparsely set long, erect, golden brown hairs and intermixed with short, sericeous,