cence as in rubidus and intermixed with yellowish to fuscous

simple pubescence.

Holotype: 9, March 23, 1921, Dundedin, Florida (W. S. Blatchley); author's collection. Paratypes: ♀, topotypic. 9, South Carolina (Mrs. A. T. Slosson).

Lepidopsallus ovatus n. sp.

Allied to miniatus but smaller, more ovate, and head broader, with antennae pale but relatively shorter; antennal segment II barely equal to width of vertex plus dorsal width

of an eve.

Q. Length 2.6 mm., width 1.55 mm. Head: width .77 mm., vertex .37 mm. Rostrum, length 1.03 mm., just attaining middle of hind coxae, yellowish, basal segment dark reddish, apical segment blackish. Antennae: segment I, length .143 mm.; II, .57 mm.; III, .32 mm.; IV, .23 mm.; uniformly pale yellowish, segment IV only slightly dusky; segment II just equal to vertex plus dorsal width of an eye. Pronotum:

length .54 mm., width at base 1.16 mm.

General coloration reddish vellow, somewhat shining, hemelytra semi-translucent, pronotum piceous but basal angles more reddish, scutellum dark with lateral margins reddish yellow; head piceous, vertex pale bordering eyes; hemelytra reddish translucent, but becoming blackish on outer apical half of corium, cuneus uniformly reddish translucent; membrane uniformly pale fuscous, veins scarcely reddish. Ventral surface, coxae, and basal half of femora, piceous, shining; tibiae, tarsi, and apical one-fourth of femora pale, tibial spines black, fuscous points barely visible at base of spines on hind pair. Clothed with silvery white, scale-like pubescence, more thickly on pleura and sides of venter, intermixed on dorsum with simple vellowish pubescence.

Holotype: \circ , May 3, 1924, Tucson, Arizona (A. A. Nichol); author's collection.

Lepidopsallus pusillus n. sp.

This species runs to olseni Kngt. in my key to the species of the genus, but is readily distinguished by the yellowish ground color darkened with fuscous and the semi-translucent aspect of the hemelytra.

9. Length 2.2 mm., width 1.03 mm. Head: width .63 mm., vertex .30 mm. Rostrum, length .85 mm., reaching to middle of hind coxae, yellowish, apex fuscous. Antennae: segment I, length .13 mm., yellowish; II, .51 mm., gradually