Canyon (E. D. Ball). MONTANA, σ , September 3, 1912, Bozeman. σ 2 \circ June 24, 1925, Williams, Arizona (A. A. Nichol).

This species was obtained by Drake and Hottes by sweeping grasses and sedges.

Ceratocapsus tricolor new species.

Allied to *biformis*, but distinguished by the genital claspers, black color with pale yellowish clavus and reddish femora.

♂. Length 5 mm., width 1.48 mm. Head: width .77 mm., vertex .31 mm.; eyes finely pubescent. Rostrum, length 1.4 mm., reaching to middle of hind coxæ, blackish. Antennæ: segment I, length .41 mm., yellowish, blackish on the constricted basal one-fourth; II, 1.6 mm., cylindrical, scarcely equal to thickness of segment I, black, slightly yellowish at base, finely yellowish pubescent; III, .98 mm., scarcely equal to thickness of segment II, black; IV, .68 mm., slightly more slender than III, black. Pronotum: length .59 mm., width at base 1.11 mm.; impunctate, scutellum likewise impunctate, but the hemelytra with fine setigerous punctures apparent. Ostiolar peritreme with dorsal lobe broadly convex, whereas *biformis* has a small convex area.

Clothed with long, erect, golden yellow simple pubescence; devoid of the silvery scale-like hairs found in *biformis*. Black, moderately shining, hemelytra more of a fuscous black, clavus pale yellowish translucent, embolium usually somewhat yellowish; membrane pale, apical half fuscous, femora reddish, becoming darker at base; coxæ pale, fuscous at base, front pair reddish to piceous; tibæ reddish brown to fuscous, tarsi yellowish, blackish apically. Genital claspers distinctive, right clasper with the ventral arm broadly arcuate, not sharply bent as in *biformis*, the dorsal arm flattened, bifid on apex.

Holotype: August 13, 1925, Mancos, Colorado (H. H. Knight); author's collection.

Paratypes: J, August 9, 1925, Veta Pass, alt. 8500 ft.; J, August 12, 1925, South Fork, Colorado (H. H. Knight). These specimens were swept from mixed growth of sedges in damp places, which seems to indicate the type of habitat.

Ceratocapsus fulvipennis new species.

Runs to *sericus* Kngt. in my key (Hem. Conn., 1923, p. 525), but differs in the smaller size, more abundant erect hairs, intermixed with fine, golden yellow, sericeous pubescence.

J. Length 4 mm., width 1.48 mm. Head: width .74 mm., width of vertex at base .326 mm., narrowest point forward .281 mm.; eyes finely pubescent. Rostrum, length 1.18 mm., reaching to middle of hind coxæ. Antennæ: segment I, length .25 mm.; II, 1.06 mm., equal