

- three times as high as wide, and black; third and fourth segments very short, fig. 20 (*Hesperophylum*, p. 19).....**Deraeocorinae**, p. 64
- Antennae with second segment cylindrical or almost so, or third segment three-fourths as long as second, fig. 92..... 7
7. Tarsal claws with only a pair of straight hairs between them, figs. 30-41, sometimes also with a small, inconspicuous, membranous area appressed to inner margin of claw, figs. 38-41.....13
- Tarsal claws with a pair of prominent, whitish, membranous lobes between them; these lobes either curved and fingerlike, figs. 54-67, or flaplike, figs. 42-53..... 8
8. Tarsal claws with membranous lobes fingerlike and arising from between bases of claws (true arolia); these either divergent, figs. 63-67, or convergent, figs. 54-62, at apex..... 9
- Tarsal claws with membranous lobes flaplike (pseudarolia), arising from inner margin of claw itself, figs. 42-53; frequently hooked to form a recess from which the pseudarolia appear to arise, fig. 52.....18
9. Arolia convergent at apex, figs. 54-62.....**Orthotylinae**, p. 74
- Arolia divergent at apex, figs. 63-67... 10
10. Pronotum with a prominent ridge running from the postero-lateral corner of the pronotum almost to the anterior corner, fig. 69; and with pleural suture situated some distance from anterior margin and terminating under ridge.....**Mirinae**, p. 124
- Pronotum with this ridge either absent or represented for only a short distance.....11
11. Posterior portion of head elongate, fig. 143, so that the eyes are situated their own length from pronotum (*Collaria*, p. 126)....**Mirinae**, p. 124
- Posterior portion of head short so that the eyes almost touch or do touch the pronotum, fig. 154.....12
12. Pronotum markedly widest at posterior margin, figs. 155, 180; hemelytra with corium and cuneus distinctly defined as sclerotized areas and set off from the apical membrane.....**Capsinae**, p. 131
- Pronotum swollen at middle, this portion as wide as or wider than hind margin, fig. 142; hemelytra with corium and cuneus merging so imperceptibly with the membrane, which is partially sclerotized, that there is no line of distinction between them; includes both macropterous and brachypterous forms.....**Mirinae**, p. 124
13. Calli greatly enlarged into a pair of broad humps occupying the anterior two-thirds of the central area of the pronotum, fig. 68....**Fulviini**, p. 61
- Calli much smaller, fig. 70; pronotum not humped anteriorly, but usually considerably humped posteriorly...14
14. Anterior margin of pronotum with a distinct, even, ringlike collar set off by a definite, deep groove, fig. 70...15
- Anterior margin of pronotum without a ringlike collar, fig. 77; at most with a flattened area, fig. 78.....17
15. Pronotum narrowed to a distinctly necklike anterior portion; head narrowed posteriorly and appearing stalked; hemelytra colorless, transparent and glassy with a Y-shaped red or fuscous mark, fig. 98 (*Hyaliodes*, p. 57).....**Dicyphinae**, p. 52
- Pronotum not greatly narrowed anteriorly, figs. 70, 71; head sometimes narrowed posteriorly but not stalked, fig. 71; hemelytra not colorless and glassy.....16
16. Eyes distinctly removed from posterior margin of head, fig. 71; pronotum with calli represented by a smooth, depressed shining area forming a second "collar," fig. 71.....**Clivinemini**, p. 64
- Eyes bordering on posterior margin of head, figs. 70, 105; pronotum with calli not depressed below level of adjacent area of pronotum.....**Deraeocorinae**, p. 64
17. Anterior portion of pronotum set off by a dark, impressed line running from antero-lateral corner to posterior margin of calli, fig. 72.....**Largideini**, p. 63
- Anterior portion of pronotum without such a line, figs. 77, 79.....**Phylinae**, p. 22
18. Dorsal outline almost circular, fig. 99, and pronotum with narrow, ringlike