

lo dilatatis, postice attenuatis, membrana unicolora, opaca, antice vix distincto punctata; corpore subtus nigro.

Long. $\frac{3}{4}$ — $\frac{4}{5}$ l. — lat. hemelytr. $\frac{1}{2}$ — $\frac{3}{5}$ l.

Des sommités du mont Patannas.

Deraeocoris rubrovulneratus Motsch.

Figura elongato-ovata Ph. *fulvomaculati* Debeer, *sed minor; punctatus, ater, thorace nitido, hoc angulis posticis, scutelli apice, axillis geniculisque rufo-testaceis, membranis margine interiori postice vulnerato, rubro, tibiis antice subalbidis; capite triangulari, glabro, nitido, fronte inter oculos maculis duabus rufis vel testaceis, oculis transversis, prominulis, fusco-cinereis; thorace transverso-trapezoidale, nitido, fortiter punctato, subconvexo; scutello triangulari, punctato-rugoso, subnitido; hemelytris basi sparsim punctatis, elongatis, medio vix dilatatis, membrana postice sinuato truncata, lateraliter incisa, apice angulatim producta.*

Long. $1\frac{1}{2}$ —2 l. — lat. $\frac{2}{3}$ — $\frac{2}{3}$ l.

Des Montagnes de Nura-Ellia.

Deraeocoris viridanus Motsch.

Figura Der. *pratensi* et Der. *icteocephali*, *sed paulo minor, oblongus, punctato-rugulosus, nitidus, melinus, capite, antennarum basi, corpore subtus pedibusque magis testaceis, oculis oblongis, prominulis, nigris, antennarum rostelloque apicis unguiculisque infuscatis; capite minuto, triangulari, vertice inter oculos transversim impresso, nitido, impunctato, antennis corporis fere longitudine, valde setu-*