

spot behind the middle and another near the apex; cuneus transparent; narrow borders of cuneous and cell of membrane dusky. Legs distinctly whitish; tarsi and claws dusky. Genital segments whitish.

Three examples from Guatemala: Gualan, Jan. 14th, 1905; Santa Lucia, Feb. 2d, 1905.

***Eccritotarsus atratus* Distant.**

A good series from Guatemala: Gualan, Jan. 14th; Santa Lucia, Feb. 2d; Los Amates, Feb. 18th to 28th, 1905.

***Eccritotarsus nocturnus* Distant.**

Two specimens, taken at Gualan, Guatemala, Jan. 14th, 1905.

***Eccritotarsus procurrens* Distant.**

A large series from Guatemala: Gualan, Jan. 14th; Los Amates, Jan. 17th, and Feb. 25th; Santa Lucia, Feb. 2d; Puerto Barrios, March 3d, 1905.

***Annona bimaculata* Distant.**

A large series from Guatemala; Los Amates, Jan. 17th, Feb. 18th to 28th; Santa Lucia, Feb. 2d, 1905.

***Annona decoloris* Distant.**

One specimen, taken at Santa Lucia, Guatemala, Feb. 2d, 1905.

***Neofurius tabascoensis* Distant.**

One specimen, taken at Mazatenango, Guatemala, Feb. 3d, 1905.

***Bibaculus modestus* Distant.**

Eight specimens from Guatemala: Santa Lucia, Feb. 2d; Puerto Barrios, March 3d, 1905. As noted by Distant, this is a variable species. Three specimens are typical *modestus*; the others have the black markings brownish and more or less indistinct.

***Neosilia pulchra* Distant.**

One specimen, taken Feb. 2d, 1905 at Santa Lucia, Guatemala.

***Neosilia viduata* Distant.**

A good series from Guatemala; Los Amates, Jan. 17th and Feb. 18th to 28th, 1905.

***Jornandes parvus* Distant.**

Two specimens from Guatemala: Los Amates, Feb. 18th and 25th, 1905. The two examples differ from Distant's description in having the outer half of the first antennal segment swollen and black.

***Lampethusa anatina* Distant.**

A large series of this variable species from Guatemala: Los Amates, Jan. 17th and Feb. 25th; Santa Lucia, Feb. 2d; Mazatenango, Feb. 3d; Puerto Barrios, March 3d, 1905. One specimen from Honduras; Feb. 21st, 1905.