

Syn.: *Cimex leucocephalus* LINN., Syst. Nat. Ed. X, 446, 46. Fn. Sv. 251, 940. DEGEER, Mém. III, 290, 28. *Lygaeus* id. Ent. Syst. IV, 175, 140. WOLFF, Ic. Cim. II, 76, 73, T. XIII, f. 73. PANZ., Fn. Germ. XCII, f. 12. *Phytocoris* id. FALL., Hem. Sv. 111, 67. HAHN, Wanz. Ins. II, 88, f. 174. *Capsus* id. H. SCH., Nom. Ent. p. 53. MEY., Rh. Schw. 109, 100. F. SAHLB., Mon. Geoc. 117, 59! FLOR, Rh. Livl. I, 358, 51. *Attus* id. BURM., Handb. II, 276, 1. *Phytocoris (Attus) leucocephala* ZETT., Ins. Lapp. 276, 32! *Strongylocoris* id. BLANCH., Hist. d. Ins. 148, 1. COSTA, Cim. Neap. III, 48, 1. REUT., Rev. synonym. n:o 257. *Capsus (Leptomerocoris)* id. KIRSCHB., Rh. Wiesb. 86, 114. *Stiphrosomus* id. FIEB., Crit. 24. *Stiphrosoma* id. FIEB., Eur. Hem. 281, 1. DOUGL. et Sc., Br. Hem. 482, 1. STÅL, Hem. Fabr. I, 88, 1. REUT. Hem. Gymn. Sc. et Fenn. 104, 1! SAUND., Syn. 288, 1. *Capsus (Lopus)* id. THOMS., Op. ent. 433, 53! *Halticus* id. VOLL., Hem. Neerl. 171. — *Cimex decrepitus* FABR., Ent. Syst. IV, 125, 178! *Miris* id. LATR. Hist. Nat. XII, 228, 33.

Var. β steganoïdes J. SAHLB.: Minor, capite toto coxisque posterioribus piceis vel obscure piceo-ferrugineis; pectore toto nigro vel medio piceo. Long. $3\frac{2}{5}$ — $4\frac{1}{2}$ mm.

Syn.: *Stiphrosoma steganoïdes* J. SAHLB., Not. Skpts pro F. Fl. Fenn. XIV, 1875, 306! REUT., Hem. Gymn. Sc. et Fenn. 104, 2! *Strongylocoris leucocephalus var.* REUT., Rev. synonym. n:o 257.

Var. γ sibiricus REUT.: Capite corpori toto concolore, nigerrimo. Long. 4—5 mm. (An species propria?).

Hab. in *Galio* (MEYER-DUER, ipse), *Campanula rotundifolia* (DUDA), *C. rapunculoide* (MONTANDON), etc. in toto territorio: Lapponia; Scandinavia; Fennia; Livonia; Rossia (Moskva, Charcov, Mohilev, Kasan, Simbirsk, Chvalynsk, Orenburg, Sarepta, Ural!); Dania; Germania tota; Britannia; Batavia; Gallia; Helvetia; Italia; Austria tota; Hungaria; Moldavia; Graecia; Asia minor; Caucasus; Algeria. *Var. steganoïdes* praecipue in albis et in parte boreali territorii inventa: Lapponia rossica (Kantalaks!, D. J. SAHLBERG, Kaschkarantsa!, D. LEVANDER); Svecia in *Galio* (Holmiac, ipse); Dania!, D. SCHLICK; Anglia!, D. Dr. PUTON; Germania (Breslau!, D. HAHN); Tirolia!, D. GREDLER; Austria inferior (Lenz in *Galio*, D. P. Löw). *Var. sibiricus* in Sibiria (Sujetuk!, Osnatjennaja!, D. HAMMARSTROEM, plura specimina).

Descr.: Caput basi pronoti multo angustius, latitudini verticis interoculari aequae longum, rubrum totum (♀) vel clypeo verticisque margine medio magis minusve piceis (♂), rarius totum rufo-piceum vel piceum vel nigrum; fusco-pilosulum, pilis in certo lumine pallidis. Rostrum coxas intermedias attingens, rubrum, articulo ultimo nigropiceo. Antennae nigrae, articulo primo vel (♀) etiam secundo basin versus rubris, secundo margini verticis interoculari aequae longo, duobus ultimis secundo parum vel paululum longioribus, quarto tertio fere $\frac{1}{3}$ brevior, apice saepe ferrugineo. Pronotum basi longitudine paulo magis quam dimidio latius, versus apicem modice angustatum, disco versus apicem sat declive, sat dense magis minusve fortiter (in minoribus fortius) impresso-punctatum, nigrum, dense longeque griseo-fusco-pubescentem. Scutellum nigrum, transversim strigosum, longe fusco-pubescentem. Hemielytra maris et feminae abdomen brevius superantia, subtilius densius punctata, longe et dense fusco-pubescentia, pilis in certa luminis directione griseis; membrana obscure fusca, macula parva ad apicem cunei