

Capsidae novae mediterraneae

descriptae

ab

O. M. Reuter.

III.

1. *Tuponia obscuriceps* Reut.

Diagn.: Virescens, superne dense albo-pubescens, capite, saepe etiam callis pronoti scutelloque basi vel fere toto in ferrugineum vergentibus; cuneo toto concolore; membrana sat fumata, venis virescentibus, areolis nebulaque infra apicem areolae minoris obscurioribus, macula ad apicem cunei hyalina; tibiis spinulis nigris e punctis distinctis nigris, maris minoribus, feminae majoribus, nascentibus; capite basi pronoti parum vel paullulum (σ^7) vel ad summum $\frac{1}{6}$ (φ) angustiore, vertice oculo circiter $\frac{4}{5}$ vel fere duplo (σ^7) vel circiter $2\frac{1}{3}$ (φ) latiore; oculis exsertis, granulatis, praecipue maris maximis, fere totis ultra angulos apicales pronoti excedentibus; antennis articulo secundo latitudine capitis paullo basique pronoti parum longiore (σ^7) vel latitudine capitis paullo longiore basique pronoti ad summum $\frac{1}{6}$ brevior (φ). Long. σ^7 $2\frac{3}{4}$ —3, φ $2\frac{1}{2}$ mm.

Hab. in *Lymaniastro Guyoniano*: Algeria (Biskra! d. 16 febr. 1894), D. E. Saunders.