

## 9. *Atomophora maculosa* Reut.

Superne albida, opaca, albo-pilosa; capite, pronoto scutelloque magis minusve dense nigro-fusco-variegatis et punctatis, maculis saepe confluentibus, vertice basique frontis medio parum signatis, interdum punctis ochraceis vel ferrugineis notatis, clypeo lineis duabus fuscis; basi scutelli maris tota nigro-fusca; clavo, corio et cuneo albidis, punctis nigro-fuscis conspersis et adhuc maculis sat magnis nigro-fuscis aequaliter adpersis, angulo cunei interiore nigro-fusco; membrana cum areolis fusco-fumata, albo-hyalino-irrorata, venis albidis, cubitali versus apicem connectenteque saepe ferrugineo-variegatis, macula sat magna ad apicem cunei aliaque plerumque majore media limbi externi albo-hyalinis; antennis ( $\sigma^{\circ}$   $\varphi$ ) virescentibus, apicem versus testaceis, articulo primo albedo, annulo anteapicali fusco; pedibus albidis, femoribus apice fusco-punctatis, anterioribus apice haud vel parum, posticis apice late infuscatis, tibiis albospinulosis, margine superiore punctis magnis nigro-fuscis signatis, inferiore innotatis, tarsis pallidis vel solum articulo tertio apicem versus infuscato; rostro apicem coxarum intermediarum attingente; vertice oculo aequo lato ( $\sigma^{\circ}$ ) vel hoc fere duplo latiore ( $\varphi$ ). Long.  $\sigma^{\circ}$   $3\frac{4}{5}$ ,  $\varphi$   $3-3\frac{1}{3}$  mm.

*Var.  $\beta$*  capite ferrugineo-signato.  $\sigma^{\circ}$ .

Hab. in Transcaspia (Ashabad! d. 19 maji 1902), D. C. Ahnger (Mus. Helsingf.).

*A. pantherinae* Reut. sat similis, differt autem colore antennarum, cuneo aliter picto, tibiis inferne late innotatis tarsisque pallidis. Caput basi pronoti circiter  $\frac{1}{3}$  ( $\sigma^{\circ}$ ) vel  $\frac{2}{5}$  ( $\varphi$ ) angustius, ab antico visum latitudini frontis oculique unici fere aequo longum ( $\sigma^{\circ}$   $\varphi$ ), a latere visum altitudini basali longitudine aequale ( $\varphi$ ) vel subaequale ( $\sigma^{\circ}$ ), clypeo prominente, angulo apicali acuto, genis angustis ( $\varphi$ ) vel haud distinguendis ( $\sigma^{\circ}$ ), gula brevi ( $\varphi$ ) vel haud distinguenda ( $\sigma^{\circ}$ ). Oculi longissime in genas extensi ( $\varphi$ ) vel gulam attingentes ( $\sigma^{\circ}$ ), nigro-fusci, granulati, orbita interiore usque ad medium paralleli, dein sinuati. Rostrum albidum, ipso apice nigro, articulo primo caput vix superante. Antennae sat longe supra apicem oculorum in sinu eorum ( $\sigma^{\circ}$   $\varphi$ ) insertae, articulo