

with the eyes subglobose and almost pedunculate; the face prominently convex, almost vertical in front, curving inferiorly; the tylus prominent; superior cheeks almost vertical, broad triangular, with the apex placed inferiorly; lower cheeks placed in large part behind the upper ones, thickened, blunt at the lower end; bucculæ slenderly lunate, concave inside, as long as the throat; rostrum short and stout, inserted on a line running direct from the base of the antennæ; occiput with a carinate collar, which is convexly elevated on the middle, and with an incised line bounding it in front; antennæ stout, about as long as the pronotum and body united, the basal joint about as long as the head, a little thickened beyond the base, second joint very long, less stout than the basal one, cylindrical, the two apical joints a little more slender. Pronotum transverse, the sides oblique, narrowing anteriorly, the anterior angles rounded, anterior margin moderately concave, with the callosities broad, bilunate, each lobe convex behind, and with the space uniting them concave behind. Femora broad, compressed; hind tibiæ very long. Scutellum triangular, the three sides almost equal, and the surface very feebly convex. Hemelytra short and wide, with the costal margin broadly accuated posteriorly, with the cuneus very large and broad, and the membrane short and attached to the inner side at the end of the corium.

*L. chloriza*, new sp.

Pale, clear-green, densely pubescent. Head yellowish; face with a blackish circle in front, which is interrupted above; each side of vertex with a dark-brown dot, and the base of tylus, the suture at the tip of lower cheeks, and the eyes dark piceous. Antennæ pale piceous, paler on the basal joint. Rostrum reaching to the intermediate coxæ, tinged with piceous at the ends of the joints and on the apex; the basal joint a little longer than the throat, the second a little longer. Pronotum narrower in front than behind, trapezoidal, the sides oblique, the posterior margin straight and the anterior one a little concave; surface green, somewhat yellow anteriorly, with a dark dot each side of posterior division, and a brown curved line bounding each lobe of the callosity posteriorly. Under side whitish-green, or very pale yellow, with a black spot at the base of the anterior coxæ, and one above; also a faint duskiness at base of the other coxæ. Legs bright yellow, or a little greenish, the tarsi somewhat infuscated at tip, and the nails piceous. Scutellum faintly tinged with yellow, closely coated with whitish hair. Corium green, unevenly punctate, the punctures shallow, and sometimes confluent, those of the claws coarser; the surface polished, but closely invested with pale, almost erect pubescence; membrane white, with a faint cloud of fuliginous outwardly. Tergum black on the disk, the broad connexivum and the apex orange, the disk smooth, and all the remaining surface invested with remote prostrate hairs, venter polished, remotely pubescent, yellowish, but a little obscured at base.