

fourth a little more slender, as short or shorter than the basal joint. Pronotum a little wider than long, broadly a little convex, the sides rounded off, the lateral margins concave; posterior margin arcuated, the anterior margin concave; callosities large, convex, transverse, bounded behind by an impressed line, separated in the middle by a transverse indentation. Legs stout, the thighs compressed, posterior tibiæ very long; basal joint of tarsi shorter than the second. Scutellum almost equilateral, the surface even, very feebly convex. Hemelytra flat, the costal margin strongly carinate, almost straight.

1. *S. chloris*, new sp.

Pale green, or greenish-white, elongate, pubescent. Head long, five-angled, almost vertical, pale green, polished, clothed with sparse, long, whitish pubescence; center of the vertex posteriorly with a short black line, indented in front, which connects with a subquadrangular black ring bounding the inflation of the cranium; face black, highly polished, the superior cheeks green; space beneath, before, and above the eyes also green; throat and base black. Eyes brown. Bucculae margined with green. Rostrum testaceous, reaching to the back of the posterior coxæ, the third and fourth joint piceous. Antennæ as long as the hemelytra including the membrane, black or piceous, with the two last joints dusky; basal joint a little shorter than the width between the eyes; second joint as long as from the front of cranium to the base of the pronotum; the third a little shorter; fourth shortest. Pronotum transverse, green, yellow anteriorly, clothed with erect, long, whitish pubescence, the middle longitudinal line a little depressed, and sometimes having a few black points anteriorly; each side of middle with a round black dot; anterior angles obliquely rounded, the posterior ones prominent, with the margin behind them arcuated. Scutellum bright green, sparsely clothed with white hairs, the basal middle with a black spearhead-mark. Sternum black; the deflected propleura green, polished; the other pleural segments blackish, margined with white. Coxæ more or less dusky, but green at tip; legs green, the tibiæ at the tip and the tarsi piceous, the nails black. Corium and clavus apple-green, sometimes a little sprinkled with black, obsolete rugulose, moderately clothed with white, long, oblique, pubescence; membrane smoky, paler at base, the nervures greenish; wings smoky. Venter black, broadly greenish-white each side of disk, neatly clothed with silky white pubescence, the segments more or less margined with white.

Length to tip of venter $3-3\frac{1}{2}$ millimeters. Width of pronotum $1\frac{1}{2}$ millimeters. Length of hemelytra $3\frac{3}{4}$ millimeters.

An elegant little Capsid, of which I found a few setting singly upon the flowers of a slender pale Composite, growing in Manitou Park, August 14. Another specimen was swept from some weed in the vicinity of Colorado Springs, in the low grounds near the Fountain Creek.

2. *S. malina*, new sp.

Larger and more slender than the preceding species, with longer an-