

areole very wide and long, the inner areole very distinct, about one-third as wide at tip as the preceding. Middle joint of tarsi very short, the basal and apical ones long, nearly equal.

B. amicta, new sp.

Ivory yellow, or pale fulvous; form much like that of *Geocoris bullatus* Say. Head smooth, an oblong spot in each angle at base of vertex, a crescent each side bounding the convexity of the face, transverse ridges of the front placed each side of a depressed longitudinal line, three spots above the tylus, middle line of the latter, and margins of all the segments of the cheeks rufous; sockets of the antennæ, basal joint of the same, and basal joint of the rostrum dark piceous. Base of vertex with an impression along the front of the carina, and this connects with the longitudinal indentation each side, and next below which is an arcuated, impressed, scabrous line bounding each side of the front; basal carina almost straight, the occipital area piceous, triangularly excavated each side; antennæ rather slender, excepting the basal joint, dusky, paler at tip, the second joint of equal thickness throughout, much thinner than the basal one, and about as long as the eye and pronotum taken together; rostrum pale piceous beyond the basal joint, reaching over the tip of the posterior coxæ ♀, but extending upon the base of the last ventral segment ♂. Pronotum polished, a little narrower in front than at base, ivory-yellow, roughly unevenly punctate, piceous on the collum, posterior margin, and on the lateral margin at base; fore part of the disk fulvous each side. Margins of the pleural segments, and more or less of the coxæ, rufous. Scutellum ivory-yellow, piceous across the base, polished, minutely wrinkled. Corium feebly polished, opaque, obsoletely punctate, margined exteriorly and interiorly and with a gradually widening curved streak running back to a wavy broad band at tip, piceous; clavus more coarsely punctate, margined on both sides with piceous; cuneus ivory-yellow, broadly piceous at tip and slenderly so on the exterior margin; membrane dusky, paler at base, the nervules of the areoles pale yellow, and the areoles dusky next the tip. Legs dull testaceous, more or less piceous on the femora and tip of tibiæ, tarsi dark piceous. Venter highly polished, the sutures, incisures, ovipositor, and genitalia rufo-piceous or rufous.

Length to tip of venter 4 mm.; to tip of membrane $4\frac{1}{2}$ —5 mm.

Mr. Bolter collected several specimens in New Mexico and kindly gave me a pair of both sexes.

Div. **PLAGIOGNATHARIA.**

Psallus delicatus, new sp.

Rosy pink or testaceous tinged with dull brown; moderately robust, polished. Head almost vertical; face convex, smooth along the middle, and each side of this with a crescent composed of short fuscous transverse lines, the middle line of vertex obsoletely impressed; tylus short, prominent, tinged with pale piceous; antennæ testaceous, dusky on the last two joints, sometimes the basal and second joints pale piceous; basal shorter than the head, the second a little longer than the pronotum; tips of the cheeks and base and tip of the rostrum generally piceous, the latter reaching upon the posterior coxæ; eyes dark brown, nearly vertical, abruptly narrowed beneath where bending upon the throat. Pronotum polished, convex, transverse, remotely very obscurely punctate; the lateral margins carinate-reflexed, oblique, hardly sinuated; the callosities generally fuscous in the surrounding depressed lines;