

Linn; Illinois, by Mr. Stromberg; Rock Island, B. D. Walsh; near Evanston, Robert Kennicott; Kansas; Wisc.; Waco, Texas, Mr. Belfrage; Washington, D. C., June 7, O. Heidemann; Fredericksburg, Va., by myself; Georgia, near Atlanta, Mr. Morrison; Maryland, Anne Arundel and Prince George counties, June 5-10, by myself. This species is very near to *H. rubicunda* Fallen, and differs from it most in the proportion of the second joint of the antennæ. It is somewhat variable in depth of colors, but it has not been found of the rosy-red color common to specimens from Europe. It is generally of a pale dull fulvous color, with a whitish powdery aspect, caused by the flat, whitish pubescence, and the vein of the areole is also generally white.

17. *Camptobrochis schwarzi*. New sp.

Ivory yellow, highly polished, above conspicuously, sparsely punctate with brown; form a little less robust than that of *C. nebulosus*. Head highly polished, convex, crossed near the base by a curved brown band, the brown color spreading down each side of face and leaving an oblong oval pale spot on the middle, each side of tylus lined with piceous, sutures of the cheeks also piceous; eyes dark brown; rostrum yellowish testaceous, piceous at tip, reaching between the middle coxæ; antennæ of normal length, and medium thickness, the basal joint usually dark at tip, the second with two dark bands, that of the apex darkest, the two apical joints united scarcely as long as the second one, infuscated except at the points of articulation. Legs honey yellow, minutely sprinkled with brown, the femora with two piceous bands near the tip, and sometimes a little piceous beneath, tibiæ also with two wider bands, the nails and tip of tarsi usually piceous. Pronotum strongly convex, rather coarsely, unevenly, roughly punctate with pale brown, the callosities prominent, convex, either all black, or spotted with black or dark brown, the collum sharply defined, ivory white, preceded by a piceous border on the head, posterior margin white, sharply defined, obsolete sinuated next the scutellum, the sides widely rounded off, sinuately bordered, with the lateral flap pale and carrying a piceous spot behind the eye. Scutellum prominently convex, highly polished, ivory yellow, with a large, black, obconical spot running out from the base to next the tip, which is set with a very few coarse punctures. Beneath pale testaceous, becoming darker on the sides and venter, the prosternum with a small black spot in front of the anterior coxæ, back of this a black spot appears farther out next the meso-pleuron and from a series extends backward to the base of the ultimate segment of the venter. Hemelytra either ivory yellow, or pale testaceous, deeply somewhat irregularly punctate, minutely pubescent, convexly inflated in the middle, the punctures of the disk and behind being generally darker brown, the coarse pale vein of the corium callous on the middle and marked by a dark brown spot which often continues back in a streak to the inner corner of the cuneus; the cuneus pale, marked at tip with a dark brown spot; the posterior border of the corium is also some-