

times brown; membrane faintly tinged with brown, vein of the basal areole interruptedly brown. Length to end of abdomen, 4-4½ millim; to tip of membrane, 5-5½ millim; width of base of pronotum, 2 millim.

This pretty insect stands between *C. nebulosus* and *C. grandis*. It is variable in the amount and depth of dark marking of the surface, and that depends to some extent upon the maturity and vigor of the individual. Soft specimens are almost always paler and less extensively marked than those which are firmer and stronger.

Several specimens, mostly males, were captured at American Fork, June 22, 1891. I have examined others which were collected by Dr. Hagen near Ellenborough, on the Yakima river, Washington Territory, July 8, 9, 1882; and others have been in my possession which were brought from British Columbia by Robert Kennicott, and from other parts of the northwest territories of the United States by various persons.

18. *Phytocoris eximius* Reuter. Ofvers. Kongl. Vetens.-Akad. Förhandl., 1875, p. 67.

Three specimens were taken in the American Fork Cañon, June 23. The species is now known to have a very extended distribution. I have examined specimens which were collected in Massachusetts, New York, New Jersey, Pennsylvania, Maryland, North Carolina, District of Columbia, Virginia, Georgia, Florida, Texas, Arizona, and Northern Mexico.

19. *Orthotylus*. New sp.

One specimen from near Great Salt Lake, June 13. Not sufficient for description.

20. *Orthotylus*. New sp.

One specimen was taken in the American Fork Cañon, June 23. It will not admit of description.

21. *Psallus*. New sp.

A single specimen was obtained in the American Fork Cañon, June 23. It will not serve for correct description.

22. *Psallus*. New sp.

One specimen was secured at Wasatch, June 27. It will not serve for description.

23. *Asciodema inconspicua*. New sp.

Greenish-white, long and narrow, tender, dull, the upper surface with minute erect, black remote pubescence. Vertex a little yellow, transversely depressed above, followed below by the curved, impressed bounding line, and a longitudinal faintly impressed line on the middle, occiput carinate-ridged, front smooth, not apparently punctate, moderately convex; rostrum reaching between the middle coxæ, piceous at tip, the tylus stout and very prominent, antennæ long, the basal and second joints stout, more or less discolored, the second fuscous at base. Pronotum transverse,