

joint paler, minutely bristly, a little longer than the clavus, the third joint scarcely more than half as long; rostrum pale fulvous, darker at tip, reaching to the posterior coxæ. Pronotum short and very convex, infuscated each side and across the base, the callosities dark brown, transverse, wide apart, the surface minutely pubescent, and the posterior edge yellow, curved, indented in the middle and next the humeral angles; the pleural flap dusky, broadly pale beneath. Anterior coxæ pale yellowish, the femora pale fulvous, a little sprinkled with brown, especially towards the tips, tibiæ paler, set with pale bristles; tarsi dark at tip including the nails. Scutellum tumid, indented at base, bright fulvous. Hemelytra bright fulvous, long, not much curved on the costal margin; cuneus of the same color; membrane yellowish white, spread with branching brown lines on the sides and tip, the looped vein deep yellow. Venter a little dusky on the disk, sometimes pale fulvous sprinkled with rufous.

Length to tip of venter, $4\frac{1}{2}$ –5 mm. Width of pronotum, 2 mm.

One specimen was taken near Cape St. Lucas by Mr. John Xanthus. In the present collection there are a few specimens marked Cal. 11, and I have examined others which were collected at San Bernardino, Los Angeles and at Flagstaff, Arizona.

COMPSOCEROCORIS ROSEUS n. sp.

More slender than the preceding species, with the outer margin of the corium nearly straight; color roseus, or pale yellow tinged with rufous. Head moderately long, very convex, finely hoary pubescent, distinctly contracted into a neck behind the eyes, the eyes large and very prominent, dark brown; rostrum slender, pale yellow, reaching over the second ventral segment. Antennæ