

*PÆCILOCAPSUS MARMORATUS* n. sp.

Pale yellow, short, moderately robust, hardly polished, very minutely scabrous, most minutely and remotely pubescent. Head polished, impunctate, nodding, strongly convex in front, with the eyes quite small, pale brown, tylus long, tapering, continuing almost the same curve as the front; antennæ moderately slender, long, rod-shaped, and very gradually tapering after the apex of the second joint, dark brown, the basal joint longer than the head, marked with several irregular or diagonal yellow bands, the second joint as long as the clavus, about uniformly thick throughout, crossed by a narrow yellow band before the middle, the third and fourth joints short, subequal, each with a narrow whitish band at base; rostrum pale yellow, very slender towards the tip, reaching beyond the posterior coxæ, a little piceous both at base and tip. Pronotum a little wider than long, convex, marbled with dark brown on the sides and posterior two-thirds, the lateral margin slenderly carinate, with a slender brown line on it and another above it, collum narrow but clearly defined, bounded before and behind by a slender brown margin, on the middle behind the callosities are two round, brown dots, humeral angles more projecting than the lateral margin, the posterior margin a very little curved, but bending down towards the scutellum; pleural flaps quite narrow, scabrous. Legs yellow, the femora flecked with brown, and the tibiæ with the knees, two bands and the tip also brown, apex of the tarsi, and the nails piceous. Scutellum moderately convex, usually marked with two longitudinal brown streaks, the tip minutely acuminate. Underside whitish yellow. Hemielytra pale luteous, spread with close golden-yellowish pubescence, irrorated and marbled with brown in such a way as to leave numerous dots and irregular spots of the surface exposed, the base