

6). Southern Colorado, July (Carpenter—see Uhler, 7). Yellow varieties on weeds near water, in suburbs of Denver. Green variety, marked with fuscous, from rank growing plants in Beaver Brook Gulch, August 6th (Uhler, 5). Ula, Custer County, July 30th (Cockerell, 10). Estes Park (Snow—see VanDuzee, 5).

Fort Collins, May 8th to October 7th and June 9th, on alfalfa and barley (Baker). Foot-hills five miles west of Fort Collins, May 11th to 19th; Manitou, June 25th (Gillette) Veta Pass, June 21st (E. A. Schwarz).

*Teratocoris discolor* Uhl.

Garland, June 18th (E. A. Schwarz—see Uhler, 3).

*Teratocoris longicornis* Uhler n. sp.

"Elongated, bright green, resembling a Miris. Head very slightly nodding, green, broad, the front tabulated above and bounded by deep sutures, black anteriorly along the depressed middle line and also each side, the vertex produced in an arch posteriorly, and with a black arc at base; tylus black, rostrum reaching midway between the middle and posterior coxae; mostly greenish-testaceous, piceous at tip; antennae long, the basal joint green, the other joints dull reddish, the basal one as long as the vertex and pronotum united, the second as long as the pronotum and clavus united, the third and fourth broken from the specimen. Pronotum trapezoidal, a little longer than wide, flat anteriorly, the sides gradually oblique, with the edge reflexed, a little constricted behind the line of the callosities and crossed there by a black band, the middle line unevenly impressed, black, connected behind with a broad black band across the base, which is obsolete punctate and wrinkled, the submargin before the humeri tabulated and with the edge not reflexed, posterior margin very slightly sinuated. Sternum whitish along the middle. Legs green, the tibiae and the tarsi more or less yellow, with the nails and apex blackish. Scutellum black, with a large yellow spot near each basal angle. Hemelytra green, long elliptical, minutely rugulose and punctate, the costal border prominently reflexed; membrane long, dull whitish, faintly dusky at base, including the base of vein. Mesosternum, posterior coxae and middle and apex of venter, pale yellowish, the abdomen minutely pubescent.

Length to tip of abdomen 4.5 mm. To tip of membrane 5.5 mm. Width of pronotum 1.25 mm. This is a well marked species of which only a single male was sent to me from Colorado. The genital hook is placed on the left side, and is long, stout and moderately curved. In this specimen the base of the scutellum is broadly exposed. A series of both sexes is greatly desired for comparative study."

Steamboat Springs, July 12th on Carex (Baker).

*Leptoterna amoena* Uhl.

Steamboat Springs, July 14th and 26th (Baker and Gillette). Estes Park, July 10th; North Park, July 20th; Colorado Springs, August 3d (Gillette).

*Resthenia atripennis* Reut.

Foot-hills twelve miles west of Fort Collins, June 30th (Gillette).

*Resthenia bivittis* Stal.

Steamboat Springs, July 26th; North Park, July 20th (Gillette).