

the second and third joints nearly of equal length. Head dull black, hairy, short, and transverse above, the vertex transversely impressed, prominently elevated behind; face tumid and convex, marked with a yellow stripe on the middle and a line of the same color on the groove next the inner margin of the eyes; cheeks, throat and bucculae pale greenish yellow; rostrum stout, greenish, black on the two apical joints, reaching upon middle coxae; antennae short, thick, the second not quite as long as the width of the base of pronotum, the third a very little shorter and not much thinner. Pronotum transverse, red or orange, set with erect black hairs, the collum white, and the band including the callosities black; lateral margins sharp, but not very prominent, pleural and sternal areas pale greenish, with the sutures and the areas between the coxae black. Legs pale fulvous at base, the apex of femora, and the tibiae and tarsi black. Scutellum pale rufous or yellow with the base blackish. Corium pale yellowish, with a large long black spot running back to tip, clavus nearly or all black, cuneus pale yellowish, bordered inwardly with black, membrane blackish. Abdomen pale greenish, with the ovipositor, genitalia and some short slender lines on the sutures of the segments black.

Length to tip of abdomen 3.5–3.75 mm. Width of pronotum 1.12–1.25 mm. Specimens of both sexes were kindly sent to me by Prof. Snow and Prof. Gillette. The first were collected at Colorado Springs, in July, by Mr. E. S. Tucker. Damaged specimens, from alcohol, were obtained for me in Dakota, and I captured a weather-worn specimen west of Denver in the month of August."

***Hadronema princeps* Uhl.**

Fort Collins, June 4th; Rist Canon, June 13th to July 30th; Gore Pass, July 29th; North Park, July 20th (Gillette). Steamboat Springs, July 12th to 16th, on *Delphinium occidentale* and other low plants (Baker).

***Hadronema pulverulenta* Uhl.**

Colorado (Gillette—see Osborn, 1).

Fort Collins, May 6th, about roots of a rose bush (R. C. Stephenson). Soldier Canon, May 19th; Golden, April 30th (Gillette). Fort Collins, September 26th, on elm (Baker).

***Hadronema robusta* Uhl.**

Colorado (Cockerell—see Ashmead in litt.).

***Pachytropis nubilus* Uhl.**

Steamboat Springs, July 26th (Gillette).

***Phytocoris colon* Say.**

Colorado (Gillette—see Osborn, 1).

***Phytocoris interspersus* Uhler n. sp.**

"In form similar to *P. eximus* Reut., but larger, and almost flat on the hemelytra. Pale delicate green, mottled with very pale brown all over the hemelytra, which are almost hirsute and the hairs interspersed with dark bristles. Head of medium size, convex, smooth, pubescent in front and on the sides; the eyes brown, prominent, reniform, placed almost vertical; front feebly grooved to near the back line of the eyes; rostrum thick at base, reaching well behind the posterior legs, dusky at tip; the antennae longer than the body, whitish, set with long bristles, the second joint longer than from the front to the posterior coxae, less hairy than the basal joint, the third a little more slender, about two-thirds the length of the second, the fourth much shorter and equally slender, the basal joint of medium thickness, about as long as the fore femur. Pronotum smooth, much wider than long, obsoletely veined with fulvous, contracted before the middle, a