

Little bristly on the sides and anteriorly, the region of the callosities transversely sunken, lateral margins deflexed, thick, fringed with dark bristles, collum narrow, whitish. Legs long, pale yellowish green, the posterior femora mottled with pale brown, tibial spines long, blackish. Scutellum convexly prominent, bald, polished, with the tip bent down, two black dots just back of it, and the lateral edge a little reflexed. Clavus and corium indented, rugose, set with silvery and dark hairs, the lateral margins gently curved, the costal submargin interrupted with pale brown lines; the membrane very large, faintly smoky, minutely flecked with brown. Venter whitish green, pubescent, polished.

Length to end of abdomen 4 mm. To tip of membrane 6 mm. Width of pronotum 2 mm. Only one specimen of this neat insect is known to me. It is a female taken in Cheyenne Canon, Colorado Springs, July (Tucker); it is somewhat above the average size of the species in this genus, and it bears a general resemblance to some of the green types of the genus *Oncotylus*, which might cause it to be confused with some of them."

***Phytocoris pallidicornis* Reut**

Colorado (Riley).

***Neurocolpus nubilus* Say**

Near Denver, August 4th (Uhler, 5). Colorado (Snow—see VanDuzee, 5). Colorado (Gillette—see Osborn, 1).

Colorado Springs, Garden of the Gods (Tucker).

***Neurocolpus inops* Say.**

On small plants in Beaver Brook Gulch, next to Clear Creek Canon, August 6th (Uhler, 5).

***Compsocerocoris annulicornis* Reut**

Colorado (Snow—see VanDuzee, 5).

***Callodemas* Uhler n. gen.**

"Having the features of a greatly elongated *Calocoris*. Head vertical, wider than the collum, having a short neck, the vertex transverse, front broad tumido-conical, with the eyes reniform, prominent, and placed below the level of the vertex, line between the front and tylus deeply cut, the tylus long, thick, subcylindric, prominently curved at base; superior cheeks quadrangular, wide, almost flat, placed a little obliquely, the base of antennae very thick, deep seated directly above the quadrangular cheeks, against the inferior third of the eyes, lowe: cheeks rhomboidal, placed nearly horizontal beneath the eye; bucculae narrow, short, not extending beyond the inner line of the eye; rostrum quite thick at base, the basal joint reaching upon the anterior coxae. Anterior border of prosternum with a narrow, double, curved fold, mesosternum tumidly elevated. Pronotum transverse, abruptly sloping in front, the front border very narrow, carrying a prominent collum which is but little wider than the base of the head. Scutellum widely exposed at base, deeply incised across the disk, the posterior division triangular and almost equilateral. Hemelytra long and narrow, the costal border almost straight, veins distinct, regular, membrane long and wide, the looped vein fusiform in contour, round at the outer end, slenderly acute at the inner end, the adjoining areole but little shorter. Abdomen almost flat above, widely curving behind the base, with the connexivum broadly reflexed."

***Callodemas laevis* Uhler n.sp.**

"Long-suboval, yellow tinged with green, bald, polished. Head highly polished, eyes dark brown, pale behind; rostrum pale testaceous, piceous on the apical joint, reaching to behind the posterior coxae: antennae filiform, very long, set with black stiff hairs, the basal joint thick, as long as from the front of the eye to the apex of the middle coxae, the second not quite as stout, longer than the corium, the other joints destroyed. Pronotum rapidly narrowing obliquely towards the front, the lateral margins sinuated, with the