

humeri rounded, lobate, the posterior margin a little sinuated on the middle, and deeply excavated inwardly from the humeri, middle sometimes with two black dots. Scutellum long triangular, acute at tip, obsolete punctate in parts near base, each side next the pronotum sometimes with a black spot. Clavus a little dusky, corium with a dusky large spot behind the middle which is triangularly forked on the posterior end, the surface remotely spread with minute silky pubescence; membrane dusky at base and apex, omitting the centre of the areoles, the veins pale yellow. Legs yellow, with the femoral and tibial spines darker; tip of tarsi and the nails piceous. Tergum black, excepting the sides and end which are greenish yellow. Venter light clear yellow.

Length to end of abdomen 8 mm. To tip of membrane 9 mm. Width of pronotum 2 mm. One specimen, a female, is in the collection from Colorado, and another from New Mexico."

Glenwood Springs, August 24th (Gillette).

*Calocoris palmeri* Uhl.

Southern Colorado (Uhler, 6).

*Calocoris rapidus* Say.

Colorado (Uhler, 1 and 6). Pueblo, August 10th (Uhler, 5). Pueblo, June (Yarrow—see Uhler, 7). Custer County, midalpine (Cockerell, 10).

Fort Collins, July 14th to August 11th; July 14th on alfalfa and *Oenothera biennis*; Steamboat Springs, July 14th, on *Solidago* and other low herbs (Baker). Rist Canon, July 14th; North Park, July 20th; Estes Park, July 12th; Steamboat Springs, July 26th (Gillette).

*Calocoris superbus* Uhl.

West Cliff, Custer County, July 27th (Cockerell, 10). Colorado (Gillette—see Osborn, 1).

Fort Collins, June 23th to August 25th, and September 27th on *Senecio douglassi* (Baker and Gillette).

*Calocoris tinctus* Uhler n. sp.

"Oblong-oval, olive brownish, moderately pubescent, mostly bald above. Head long nutant, contracted obliquely on the front, the vertex a little sunken, longitudinally incised, bounded behind by a prominent ivory white carina, the front most compressed next the line of the tylus, tylus smooth, dark brown; rostrum reaching behind the posterior coxae, pale rufo-piceous; antennae long and slender, as long as pronotum and corium united, pale dull fulvous, a little darker on the last two joints and tip of second joint, the second joint a little thicker than the following ones, as long as the head and pronotum united, the third and fourth filiform, together a little shorter than the second. Pronotum wider than long, trapezoidal, moderately convex, almost regularly punctate, with the collum, lateral and posterior margins almost straight, white, the submargin and callosities black, anterior angles a little rounded; pleural flaps deep, black, polished, coarsely remotely punctate and wrinkled, sternum blackish brown, the borders of pleural areas white. Legs brown, paler at base and on the trochanters, nails black. Scutellum moderately convex, brown, pale at tip, minutely wrinkled and obsolete punctate. Clavus and corium covered with close shallow punctures, subcostal area with a broad and long pale almost hyaline stripe; membrane dusky, with a large whitish spot near the base, the veins dark; wings dusky. Venter purplish black, opaque, pale on the middle, fulvous on the last segment, all the incisures margined with white, and with a series of pale dots on the lateral submargin, pubescence hardly distinct.

Length to end of abdomen 8 mm. To tip of membrane 7 mm. Width of pronotum 2 mm.