

Only one specimen, a female, was taken in Colorado. It is peculiar from the conical and compressed form of the clypeus, and the genital slit for the ovipositor extends back in a pale line to next base of venter. The male is a great desideratum."

Estes Park, July 15th, on Pinus (Gillette).

***Megacoelum fasciatum* Uhl.**

Near Manitou, August 13th (Uhler, 5). Manitou, July 16th (Packard—see Uhler, 5).

***Lygus annexus* Uhl.**

Colorado (Uhler, 1). Foot-hills (Carpenter—see Uhler, 6). Pueblo, July ('Wilkin—see Uhler, 7). Several var. in Beaver Brook Gulch. Pale varieties common near Denver on *Euphorbia marginata* (Uhler, 5). Denver, July 28th; Idaho Springs, July 6th (Packard—see Uhler, 5). Custer County, midalpine (Cockerell, 10).

Gore Pass, July 29th; Montrose. June 24th (Gillette). La Veta, July 4th (E. A. Schwarz).

***Lygus diffusus* Uhl.**

Custer County, high alpine (Cockerell, 10).

***Lygus guttatipes* Uhler n. sp.**

"Form of *L. prasinus* Reut., but having a narrower head, the femora on the sides and the tibial spines at base marked with black dots. Ovale, convex, dull pale green, in some specimens obsoletely tinged with brownish around the base of membrane, obsoletely scabrous and effaced punctate, set with silky stiff pubescence which is mixed with dark bristles on the sides of the pronotum. Head polished, strongly convex; eyes large, brown; antennae green, stout, the basal joint thick. a little brownish, somewhat longer than from the eye to tip of tylus, second joint growing gradually thicker towards the tip, much shorter than the base of pronotum; rostrum greenish, broadly black at tip, reaching upon the middle coxae. Pronotum transverse, convex, remotely pubescent, unevenly and finely punctate, lateral margins acute, slightly curved, the pleural piece beneath them short, sunken, having a few punctures on the middle. Legs green, the femora stout, marked with round, black spots; tibial spines black, and with a black dot at the base of each, apex of tibiae and of tarsi also black. Scutellum moderately convex, a little wrinkled. Corium and clavus obsoletely, rugose-punctate; cuneus with remote dusky pubescence, the tip sometimes minutely fuscous; membrane moderately long, pale greenish, with paler veins. Abdomen brighter green, minutely pubescent, the genital pieces blackish,

Length to end of abdomen, male 3.25, female 3.5 mm; to tip of membrane 4 mm. Width of pronotum 1.5 mm. Four specimens have been sent to me by Prof. Snow for examination. They were collected at Manitou, in August, by Mr. E. S. Tucker."

***Lygus invitus* Say.**

Steamboat Springs, July 16th (Baker).

***Lygus plagiatus* Uhler n. sp.**

"Robust, dull fulvo-griseus, or griseo-estaceous with fuscous obscure spots, opaque, pubescent. Head olivaceo-testaceous, polished, vertex incised on the middle, indented next the occipital carina, eyes brown, placed perpendicular, front convex, a little obsolete punctate, indistinctly marked with about four brown uneven stripes, cheeks and tylus brown below; rostrum flavo-testaceous, piceous at tip, reaching to the middle coxae, antennae short and moderately stout, the basal joint dark brown, the second joint a little longer than the pronotum, darker and a little thicker towards the tip, the third