

and Gillette). Manitou, September 29th on Salix; Glenwood Springs, August 24th (Gillette).

*Camptobrochis robustus* Uhler n. sp.

"Short and thick, dusky testaceous, strongly marked with fuscous and black, coarsely, and in part densely punctate. Head almost vertical, vertex short, transversely grooved, bordered with a broken fulvous line in front, the occipital carina high, fitting into the collum, ivory yellow; front bordered with pale dull yellow, polished, remotely minutely obsolete-punctate and wrinkled, closely freckled with black, the inner border of the eyes also pale dull yellow, the lower part of tylus and the bucculae yellow; rostrum pale at base, piceous at tip, reaching to the middle coxae; antennae long and slender, as long as the corium and cuneus united, mostly pale fuscous, the basal joint dull pale fulvous, obscured with fuscous, the second very long, a little stouter than the third and fourth, slightly thicker towards the tip, the third and fourth together a little longer than the second, the fourth much the shortest. Pronotum convex, coarsely, deeply, irregularly punctate in somewhat transverse wavy lines, with about four obscure stripes which widen posteriorly, the lateral margins a little curved, the humeral angles broadly rounded, the posterior margin feebly curved and sinuated, and the anterior margin contracted and bordered with a somewhat pale cellum, the callosities tumid, black, polished. Scutellum dark brown, closely and roughly wrinkled and unevenly punctate, convex, olive-fulvous at tip. Legs pale olive-brownish, the femora piceous, rough at a few points, having one or more yellowish dots near the tip, the tibiae spotted with dark brown, closely pale pubescent, with the spines, tip of tarsi and nails dark piceous. Clavus coarsely punctate and wrinkled, dark olivaceo-fuscous, corium a little paler, smoother, more finely punctate, almost bald, with the surface near the costa translucent, punctate with brown, the costal border dark brown, ending in a darker spot before the cuneus, the embolium broad and piceous black, the cuneus dark brown, bordered all around with pale testaceous; the membrane whitish, a little stained with brown at base and tip, and the veins mostly brown. Abdomen olivaceo-testaceous, finely pubescent, dusky at tip, with a line of black marks on the outer submargin, and a series of yellow dots on the connexivum.

Length to end of abdomen 5 mm. To tip of membrane 6 mm. Width of pronotum 2.5 mm. Three or four specimens have been brought to my notice. One specimen from Colorado is chiefly dark fulvous, others were mostly chestnut brown or nearly black. The pale stripe with black arrest at the end of costal area will go far towards quickly distinguishing this species."

North Park, July 20th (Gillette), and July 10th on *Artemisia tridentata* (Baker). Leadville, August 23d (Gillette). Cameron Pass, at 12,000 feet, on Salix (Baker).

*Orthops scutellatus* Uhler.

From bushes in Clear Creek Canon, August 7th. Not rare (Uhler, 5). Colorado (Uhler, 8).

Steamboat Springs, July 15th to 26th (Baker and Gillette). North Park, July 20th; Dolores, June 18th (Gillette).

*Thyriillus brachycerus* Uhl.

Colorado (Uhler, 6). Near West Cliff, Custer County, end of July (Cockrell, 10).

Estes Park, July 10th (Gillette).

*Thyriillus pacificus* Uhl.

Weld County (Uhler, 1).