

*Tinicephalus simplex* Uhl.

Colorado (Uhler 1). Foot-hills, July (Carpenter—see Uhler. 6). Mountains west of Denver, July and August (Uhler, 5).

Garland, June 30th (E. A. Schwarz).

*Pilophorus gracilis* Uhler n. sp.

"Narrower than usual, dark piceous, or nearly black, the pronotum longer than wide, and the corium distended and almost hyaline at the tip. Head depressed subconical, the vertex transversely depressed, with a central carina running back to the thick carina bounding the occiput behind; face convex, piceous, polished, becoming fulvous below; tylus narrow, pale rufo-fulvous like the cheeks; rostrum dark honey-yellow, darker at tip, reaching upon the middle coxae; antennae moderately long and slender, the basal joint short, dull fulvous, the second a little longer than the pronotum, of the same color, but darker on the gradually thickening tip, the apical joints more slender, but not abruptly decreasing, the two united scarcely longer than the second. Pronotum highly polished, semi-cylindric, with the sides a little sinuated, almost of equal width throughout, dark piceous, minutely wrinkled on the disk and behind, with the humeral angles acute and the anterior ones rounded off. Scutellum depressed behind the middle, black, and minutely scabrous. Hemelytra long, dark brown, sometimes almost black, minutely pubescent, velvety, not banded, distinctly sinuated on the sides, the membrane a little dusky, with a dark brown spot near the base, the inner border of apex of corium with a curved callous line. Legs pale dull piceous, paler on the tibiae. Abdomen black, highly polished.

Length to end of abdomen 3.5 mm. Width of pronotum 1.25 mm. One specimen, a female, lacking the tarsal joints and two apical joints of the antennae, was sent to me from Colorado. It lives on *Pinus inops* in summer, June to September, in Maryland, Virginia, New Jersey; and is also found in Massachusetts. Maturely colored specimens do not have the reddish clypeus and cheeks observed in the specimens from Colorado."

Colorado Springs, August 3d (Gillette).

*Globiceps angustata* Uhler n. sp.

"Black, highly polished, delicate, slender. Head transverse above, convex, having a few remote punctures, the eyes large and prominent, much elevated above the line of the head, the face nearly vertical, with the front a little convexly prominent, the tylus moderately narrow; rostrum very stout at base, with the basal joint long, stout and blackish-piceous, the middle testaceous and the tip piceous; antennae long, slender, dull black, the basal joint longer than the head, testaceous at base, the second barely thicker at tip, nearly as long as the costa and cuneus united, the third a little more slender, about half as long as the second, the fourth broken off. Pronotum sub-campanulate, highly polished, about twice as long as the eye; the collum contracted, narrow; the callosities very large, tumid, with a groove between, which is partly occupied by a blunt carina that runs back to posterior lobe; posterior lobe wide, convex, transversely wrinkled, with the anterior slope high, the sides steep, oblique, and gently sinuated, the humeri a little turned up and acute. Scutellum high, almost flat above, with steeply sloping sides, acute at tip. Legs pale fulvous, whitish at base, the ends of tarsi piceous. Clavus dusky, corium whitish-yellow at base, dusky on the remaining two-thirds; the cuneus pale yellowish, dusky at tip; membrane long and wide, pale fuliginous, white at the extreme base, with the looped vein dark brown. Abdomen black, highly polished.

Length to end of abdomen 2.75 mm. To tip of membrane 4 mm. Width of pronotum .87 mm. Described from a single male specimen from Colorado. It is closely related to, if not the same species as one which I found in the Province of Quebec. The female is earnestly desired for study, to complete the description of this species. This insect might readily be mistaken for one of the small Chalcididae which have black bodies and yellow legs."

Steamboat Springs, July 12th (Baker).