

Fort Collins, June 24th; Montrose, June 24th (Gillette).

*Diommatius congrex* Uhl.

Steamboat Springs, July 16th, on willow (Baker).

*Tuponia subnitida* Uhler n. sp.

"Whitish-green, long oval, minutely pubescent, with the costal margin of the hemelytra very feebly curved. Head broad and very convex, pressed back upon the thorax, smooth, not punctate; front nearly vertical, highly polished; the tylus short, compressed, but little prominent, curved beneath; the rostrum pale fulvous, reaching over the middle coxae, with the tip piceous; antennae long and slender, the basal joint short, blackish, second joint green, dark at base and tip, not much thicker than the following joint, about as long as from the front of the eye to the base of pronotum, the third a little shorter and slightly more slender, the fourth scarcely more than one-third the length of the third, equally slender. Pronotum transverse, uneven, steep, very slightly convex, smooth, impunctate; with the callosities forming an arc, preceded by a semicircular area touching the front margin; the lateral margins very oblique, sharp-edged and deflexed; the posterior margin sinuated at the base of the scutellum; the humeral angles broad and rounded. Scutellum very moderately convex, the base uncovered, the disk a little wrinkled and the tip acute. Legs greenish-white, the knees, tibial spines, dots at intervals, and the tips of tarsi black. Corium and clavus greenish-white, very minutely scabrous, and remotely effaced-punctate; membrane a little dusky behind the middle to the tip. Abdomen pale green, minutely white pubescent.

Length to end of abdomen 2.5 mm. To tip of membrane 3 mm. Width of pronotum 1.12 mm. Described from two specimens from Colorado."

Steamboat Springs, July 12th (Baker).

*Stiphrosoma atrata* Uhl.

Fort Collins, July 14th on alfalfa; Steamboat Springs, July 12th on *Carex* and *Artemisia tridentata* (Baker).

*Stiphrosoma croceipes* Uhl.

Big Narrows of Poudre river, Larimer County, July 9th on *Bigelovia* (Baker).

*Stiphrosoma robusta* Uhler n. sp.

"Black, polished, shorter, thicker, and more compact than *S. stygica* Say, with a more convex front. Head thick, convex from the vertex to end of upper cheek, the vertex with a low carina behind against the pronotum, and marked with a shallow impression each side, front highly polished impunctate, tylus narrowing a little towards the tip, deep seated in the cheeks; rostrum stout, a little piceous, reaching to the middle coxae; antennae slender, black, about as long as the head, pronotum and scutellum united, the second joint nearly the same length as the clavus, a little thicker towards the tip, the third more slender, nearly two-thirds the length of the second, the fourth about one-half the length of the second. Pronotum pubescent on the sides, with the lateral margins oblique and more bent down than in *S. stygica*, the surface scabrous, obsoletely punctate and transversely wrinkled, the callosities large, convex placed diagonally, and separated by a wide depression, the disk convex posteriorly, with the humeri acutely prominent. Pleura a little wrinkled and with a few punctures. Scutellum short, scabrous, a little punctate. Legs with stiff spines on both femora and tibiae. Clavus coarsely, closely scabrous, and obsoletely punctate, the corium a little less coarsely scabrous, and punctate, but with these features almost obliterated behind the middle, the sides and margins pubescent; membrane moderately short, brown. Abdomen broad ovate, highly polished, scarcely punctate.

Length to end of abdomen 4 mm. Width of pronotum 1.75 mm. Described from two specimens, both females, sent to me from Colorado. It inhabits also Texas and New Mexico."

Steamboat Springs, July 26th (Gillette).