

***Stiphrosoma stygica* Say.**

West Cliff, Custer County, July 2:th; Ula, July 30th (Cockerell, 10).

Steamboat Springs, July 26th (Gillette). Garland, June 30th (E. A. Schwarz).

***Halticus bractatus* Say.**

Manitou, July 24th; Colorado Springs, August 3d (Gillette).

***Labopidea chloriza* Uhl.**

Howe's Gulch, June 14th; North Park, July 20th (Gillette). Steamboat Springs, July 13th and 26th (Baker and Gillette). Aspen (W. W. Willard).

***Labops hesperius* Uhl.**

Colorado (Uhler, 1). Foot-hills and mountains, July and September (Carpenter—see Uhler, 6). From bushes growing among rocks on mountainside near Gray's Peak, August 6th (Uhler, 5).

Estes Park, July 12th (Gillette). Colorado, June, on pine (Riley). Veta Pass, July 1st (E. A. Schwarz).

***Dicyphus californicus* Stal.**

North Park, July 20th; Montrose, June 24th; Colorado Springs, August 3d; Manitou, September 29th on oak and Salix (Gillette). Steamboat Springs, July 16th and 26th (Baker and Gillette). Fort Collins. May 20th (Baker).

***Dicyphus californicus* Stal. var. *agilis* Uhl.**

Beaver Brook Gulch, August 7th (Uhler, 5).

Steamboat Springs, July 16th (Baker).

***Dicyphus vestitus* Uhler n. sp.**

"Resembling *D. famelicus* Uhler, but much less elongated, smaller, having nearly linear blackish antennae, with the apex of the second joint not thickened, and the colors darker, with a black head and mostly black scutellum. Head subglobose, a little fringed with pubescence, black, polished, with a pale spot on the vertex; rostrum dull testaceous, darker at tip, reaching to the middle coxae; antennae moderately stout, blackish, with the first joint pale towards the base, the second joint about as long as the pronotum, the third a little shorter, and not much thinner than the second, the fourth short, acutely tapering. Pronotum a little longer than wide, subcylindric anteriorly, with the sides sinuated and reflexed; the collum narrow and white; the surface highly polished, dark brown, with the middle broadly orange, which is replaced by dull white on the posterior segment; the humeri are sometimes also pale orange; callosities distinct, prominent, followed exteriorly by a swelling of the sides; posterior lobe wide, broadly sinuated behind. Base of scutellum wide, uncovered, black, the scutellum proper black with a yellow spot on each side at base. Coxae white, legs dull yellowish, with the tip of tarsi picuous. Clavus pale, with the suture dusky, corium long and narrow, obscure whitish, translucent, with a small brownish spot near the base, a dusky streak on the middle, a line of same color on the suture, and a double lobed spot of brown on the tip; cuneus white, bordered more or less with brown at tip; membrane