

*Macrotylus moerens* Uhl.

Steamboat Springs, July 16th and 26th (Baker and Gillette). Spring Canon, June 30th; Estes Park, July 10th; North Park, July 20th; Manitou, June 25th (Gillette).

*Macrotylus verticalis* Uhl.

Montrose, June 24th (Gillette).

*Bolteria picta* Uhl.

North Park, July 20th; Leadville, August 23d (Gillette). On Grizzly Creek, in mountains south-west of North Park, July 10th, on *Artemisia tridentata*; Steamboat Springs, July 12th (Baker).

*Episcopus ornatus* Reut.

Colorado Springs, August 3d (Gillette).

*Plagiognathus annulatus* Uhler n. sp.

"Closely related to *P. arbustorum* F., but narrower and smaller, brownish-black, long-oval, polished, with pale yellow legs, having the femora streaked and the tibiae annulated or spotted with black. Head longer than wide, acute in front, convex, highly polished, faintly tinged with brown; vertex concurrently convex with the front; the occiput arched and with the edge obscure fulvous; tylus scarcely more prominent than the front, curved, highly polished; rostrum reaching to the middle coxae, yellow, with the basal joint black and the apex piceous; antennae of medium length, the basal joint black, scarcely longer than the head, the second yellow in the middle, or nearly all yellow but the base, a little longer than from the front of the eye to the base of pronotum, third and fourth more slender, yellow. Pronotum deep black, highly polished, growing much narrower anteriorly, minutely, irregularly and obsoletely rugose, more convex posteriorly; the callosities large, very slightly prominent, with an indented space between, and a wavy impressed line forming the boundary between them; the lateral margins oblique, barely sinuated with the edge a little keeled, deflexed; anterior angles a little rounded, callous; posterior margin gently arcuated; the humeri prominent, almost acute, with an impression bounding them inwardly. Scutellum feebly convex, a little unevenly rugose, the tip acute and set off by a transverse series of punctures placed a little way back. Corium deep black, long, highly polished, obsoletely and irregularly rugose over most of the surface, with some punctures on the depressed borders; the clavus obsoletely and coarsely rugose, with series of shallow punctures on the sutures; base of cuneus marked by an obscurely pale sutural line; the outer angle of base of membrane marked by a triangular white spot, remainder of membrane fuliginous black. Legs pale yellow, the base of anterior coxae, the knees, rings and spines of tibiae and the tarsi black.

Length to end of abdomen 3.5 mm. To tip of membrane 4 mm. Width of pronotum 1 full mm. Described from one female specimen sent to me from Colorado. It inhabits also northern Illinois, Canada west, and New England."

Steamboat Springs, July 16th (Baker).

*Plagiognathus fraternus* Uhler n. sp.

"Larger and more flat above than *P. obscurus* Uhler, color black, minutely sericeous pubescent. Head triangular, a little longer than wide, acute at tip, with the sides a little sinuated; occipital carina indistinct, pale yellow, front convex, a little roughened on the middle; the tylus prominent, highly polished, comparatively narrow, a little curved; rostrum reaching upon the middle coxae, basal joint stout, a little longer than the throat, blackish-piceous, the middle joints fulvo-piceous, the apical joint blackish; antennae