

*Mauroidactylus angulatus* Uhler n. sp.

"Mostly pale obscure testaceous above and black beneath, long oval, minutely sericeous pubescent. Head nearly triangular, wider than long, dull black, irregularly testaceous at base, face almost vertical, minutely, sparsely pubescent, feebly convex, not apparently punctate, the occipital edge sharp, nearly straight, not carinate; the tylus narrow, almost acute at tip; rostrum yellow from base to middle, black from thence to tip, reaching to behind the middle coxae, the basal joint reaching to the end of the arched mucro, swollen at tip; the bucculae slender and pale yellow; antennae blackish, moderately stout, reaching as far as to the middle of the venter, the basal joint projecting little beyond the tip of the head, the second joint stout, of uniform thickness throughout, a little longer than the vertex and pronotum, the following joints shorter and more slender. Pronotum short, trapezoidal, transverse, a little polished, impunctate, minutely and remotely pubescent, pale greenish testaceous, almost flat, with an abbreviated black line crossing the line of the callosities, the anterior and posterior margins almost straight, the lateral margins a very little sinuated, the pleural flaps deep, obscure black, bordered with testaceous; pleural and sternal areas black, also partly margined with whitish. Legs including the coxae, greenish yellow, the middle of femora, the spines, and apex of tarsi black. Scutellum dull black, uncovered at base, and with a pale fulvous spot in each basal angle. Hemelytra obscure testaceous, minutely pubescent, placed almost flat, with the costal border a very little curved; clavus dusky towards the tip, corium with a double, long, dusky, widening spot running back to tip, cuneus with a dusky spot near tip; membrane long, pale dusky, darker in the basal cell. Venter polished, black, narrow, with the edges of the segments obscure whitish.

Legth to end of venter 3 mm. To tip of membrane 3.5 mm. Width of pronotum 1 mm. Only one specimen has thus far been brought to my notice. It is a male, and bears some resemblance to *Mauroidactylus alutaceus* Fleber., of Europe."

Steamboat Springs, July 12th (Baker).

*Mauroidactylus consors* Uhler n. sp.

"Form narrow, almost parallel sided on the hemelytra, very slightly pubescent above, black, polished. Head triangular, wider than long, acute at tip; the occipital edge sharp, a little raised at the middle; the face moderately convex, not apparently punctate; eyes prominent, large, placed nearly vertical, set close to the pronotum, dark brown; the tylus prominent, curving beneath; rostrum fulvo-piceous, darker at base and tip, reaching upon the middle coxae; the bucculae pale, narrow; antennae black, long, with the two basal joints stout, the second as long as from the scutellum to inner angle of the corium, the third a little more slender, about three-fourths as long as the second, the fourth scarcely more slender than the third, hardly one-half the length of the third. Pronotum small, moderately convex, a little wider than long, minutely scabrous and a little wrinkled; the anterior lobe short, tumidly elevated, with the impressed line feebly defined; the lateral margins oblique, hardly sinuated; the anterior margin but very little wider than the space between the eyes, with the collum obsolete, and a slight indentation behind its middle; the forward slope of surface moderately steep; humeral angles wrinkled, a little callous and acute. Scutellum widely exposed at base, longer than wide, obsolete and remotely scabrous, acute at tip. Femora broad, black, yellow at tip, the anterior and middle parts. Tibia and the tarsi pale piceous, with the spines black. Clavus obsolete scabrous; the corium a little more finely so; cuneus separated from the corium by a slender angular pale line; membrane dusky, iridescent, with the base next the cuneus and the looped vein pale. Venter black, short, polished.

Length to end of abdomen 3 mm. To tip of hemelytra 4.5 mm. Width of pronotum 1.25 mm. Only one specimen, a male with long, curved, acute genital hooks, has been sent to me from Colorado"

Leadville, August 23d (Gillette).

*Agalliastes apitatus* Uhler n. sp.

"Long-oval, polished, dark brown or black, minutely pubescent. Head short, moderately wide, highly polished, with an indented line on the middle, the vertex prominent, and a little triangular at base; front moderately convex, nearly vertical; tylus short,