

narrow, prominent, black; rostrum slender, testaceous, piceous at tip, reaching to the posterior coxae; antennae dull black, of medium length, the second joint nearly as long as the line from the tip of scutellum to the apex of clavus, the third joint a little shorter and slightly more slender, tapering, the fourth scarcely more than one-third the length of the third, still more slender. Pronotum transverse, polished, minutely pubescent, with a few sparse punctures on a line in front of the low callosities, lateral margins short, oblique, with the edge a little turned down, the collum confined to the middle of the margin, and fading into the raised surface before reaching the line of the eyes; humeral angles prominent, almost acutely callous; the posterior margin almost straight. Scutellum short, almost flat, clothed with pale pubescence, the base usually exposed, sometimes disclosing the yellowish spot at each basal angle. Legs dull yellow, the posterior pair having the femora black, all the tibiae with black spines and dots at base of spines; tarsi piceous at tip. Hemelytra covered with short yellowish pubescence, the clavus wide smoky fuscous; corium with a large fuscous, wedge-shaped spot, closed on the middle, the outer and inner borders and base and tip dull testaceous; the cuneus dusky, bordered all around with dull testaceous; membrane dusky, with the veins, base and sometimes the outer border pale. Abdomen black, polished, with a large greenish spot near the base, and some greenish spots on the pleura.

Length to end of abdomen 2.25-2.5 mm. Width of pronotum 1-1.25 mm. This species inhabits both Colorado and Kansas. It is no doubt quite common, but hitherto only a few less mature specimens have been sent to me for examination. It closely resembles *A. apicalis* Uhler, of the Atlantic States, but is a more robust and clumsy style of insect."

Fort Collins, June 4th; Manitou, June 25th (Gillette).  
Steamboat Springs, July 12th and 26th (Baker and Gillette).

*Agalliaestes associatus* Uhl.

Colorado (Uhler, 6). Pueblo (Yarrow--see Uhler, 7). Various parts of Colorado (Uhler, 9).

Fort Collins, July 24th to August 18th; Steamboat Springs, July 12th (Baker). Horsetooth Gulch, May 18th; North Park, July 20th; Trinidad, May 14th; Montrose, June 24th; Glenwood Springs, August 24th (Gillette). Colorado Springs, July (Tucker).

*Agalliaestes decolor* Uhl.

On Redstone Creek, twelve miles south-west of Fort Collins, August 1st (Baker).

*Agalliaestes fumidus* Uhler n. sp.

"Oblong-oval, pale fusco-olivaceous, clouded with smoke brown. Head narrow, polished, almost vertical, black upon the clypeus and tylus, pale greenish above, vertex strongly convex; rostrum pale towards the base, slender, reaching to the middle coxae; antennae dusky testaceous, almost piceous at base, slender, about three-fourths the length of the body, the basal joint short, piceous black, the second fuscous, nearly as long as the head and pronotum united, pubescent, the third and fourth more slender, pale, much shorter than the second, eyes prominent, dark brown, placed diagonally. Pronotum trapezoidal, transverse, bald, polished, moderately convex, pale olivaceo-testaceous, not distinctly punctate, tinged with fuliginous each side anteriorly, the collum lower than the base of the head, a little curved, with a black callous curved transverse line behind its middle; lateral margins a little sinuated, slightly notched at the outer end of the transverse impressed line, the lateral margin acute and reflexed throughout, the humeral angles moderately prominent, pale; posterior margin a little curved. Exposed base of scutellum black in the middle, yellow on either side, the scutellum proper tumid in the middle, sunken and dark at base, pale on the acute apex. Legs smoky testaceous, dark