

apical margins of the corium; margins of the cuneus, especially the outer, and the costal and subcostal nervures of the corium toward their apex thickened and colored; commissural nervure of the clavus with a blackish line at apex and sometimes at base. Nervures of the membrane from near their base to the apex of the smaller areole thick and fuscous, thus forming a double brown hook. On the disk of the corium at its widest part is a wide subcrescentic dusky cloud and a round one occupies the middle of the membrane beyond the areoles. In all my specimens there is a bluish-green mark in the larger areole next the thickened nervure and another on the base of the cuneus which may extend on to its inner margin. Wings pellucid white; tarsi brownish at apex; tip of the rostrum black.

Described from one female and three male examples in only one of which is the pale pubescence, characteristic of the genus, retained. These specimens were all taken at Leona Heights, Alameda Co., California, in August, by Mr. J. C. Bradley. This species has a little the aspect of a *Cyrtopeltis* but the different shape of the head and want of a pronotal collar will at once separate it from that genus.

***Diaphnidia capitata* n. sp.**

Smaller than *pellucida* and distinguished by its black head and fuscous basal joint of the antennæ. Length 3 mm.

Head polished, convex, the median sulcus nearly obsolete. Pronotum proportionately broader than in *pellucida*, the callousities prominent but small. Elytra parallel, scarcely widened apically, surface finely pale-pubescent; rostrum passing the hind coxæ.

Color whitish, tinged with testaceous or greenish in immature specimens. Head piceous-black, the eyes dark castaneous; antennæ a little infuscated at apex; basal joint and apex of the second infuscated in one immature specimen. Rostrum whitish, its apex scarcely darker in my specimens. Elytra diaphanous, the nervures scarcely thickened or pigmented. Membrane iridescent and very slightly enfumed beyond the areoles. Claws concolorous.

Described from three female examples; one taken by me in August 1907 at Lawton's Station, near Buffalo, N. Y., and two from "Psocid Glen" at Johnstown, Fulton Co., N. Y., taken by Mr. C. P. Alexander on August 30th 1910. The small size and black head will at once distinguish this species.

***Calocoris uhleri* n. n.**

In 1895 Dr. Uhler described a *Calocoris tinctus* in Gillette and Baker's Hemiptera of Colorado, p. 34. This name however was used by Distant in 1884 for another species in the same genus so I now change that of Dr. Uhler as above. In the Canadian