

obtuse apex incurved; sinistral finger-like or a little thickened toward its apex, a little shorter than the dextral.

Color black, polished on the front and clypeus; vertex, occipital margins and antennal sockets obscurely fulvous. Disk of the posterior lobe of the pronotum in the female broadly dull fulvous, the callosities tinged with the same color. In the male this fulvous disk is much reduced. Coxæ and legs pale fulvous, the apex of the femora and tibiæ becoming infuscated, in the male almost black, the tip of the tarsi black. Disk of the corium in the female with a whitish vitta, broadest near the base and expanded on the cuneus so as to cover all but the inner angle and apical margin. In the male this pale ray is much reduced and confined to the base of the corium. Membrane black with a pale line along the apical margin of the cuneus, which sometimes encroaches upon the apex of the nervures.

Described from one male and three female examples taken on willow at Batavia, N. Y., July 5, 1914, by Mr. H. H. Knight, who has kindly sent them to me for study and to whom it gives me pleasure to dedicate this interesting species.

Type, male, and allotype in collection of H. H. Knight. Paratype in author's collection.

30. *Orthotylus ornatus*, new species.

Size of *formosus*, but proportionately broader. Pale greenish varied with fuscous and sometimes tinged with sanguinous; median line of the pronotum and apical field of the scutellum in the female pale. Length 6mm. to tip of the membrane.

Head vertical. Vertex flat, but little depressed; the fovæ obscure, basal carina distinct. Front scarcely convex; clypeus narrow, moderately prominent. Eyes large, about two-thirds the width of the vertex; viewed from the side broad ovate, but little narrowed below, reaching nearly to the gula. Pronotum broad, rather flat; sides a little concavely arcuated, the humeri prominent; anterior angles but little rounded; surface opaque and distinctly shagreened; callosities little elevated. Antennæ short; first joint as long as the head; second nearly as long as the head and pronotum together; third and fourth conjointly about two-thirds the length of the second. Basal lobe of the scutellum somewhat exposed. Elytra broad, the costa a little