

Cuneus pale. Membrane infuscated with a short blackish ray at apex of the areole, the nervures yellowish. Antennæ black with pale incisures; the feet infuscated with the base of the femora pale. I took numbers of this neat little species about the lower end of Fallen Leaf Lake, near Tahoe, California.

7. *Macrotylus infuscatus* n. sp.

Allied to the preceding; dull fuscous with the costal margin pale or largely pale, sometimes greenish-testaceous in the female. Length 5 mm.

Head subhorizontal. Vertex scarcely flattened, the front becoming strongly convex anteriorly as in the typical forms of this genus, with a deep constriction at the base of the clypeus. Clypeus prominent, a little produced at apex as in many of our species. Eyes large, viewed from the side nearly vertical and reaching much below the middle of the side of the head. Antennæ rather long; the basal joint surpassing the apex of the clypeus by one-half its length; second two and one-half times the length of the first; the third about equal to the second; fourth flattened and about equal to the first. Pronotum long, strongly narrowed anteriorly, scarcely half as wide before as behind; sides sharply carinate throughout, a little sinuated, the humeral angles prominent. Elytra parallel or slightly widened behind; wing cell with a hamus.

Ground color pale testaceous; tinged with greenish on the pronotum, head and beneath; shaded with fuscous, especially in the males; pronotum, excepting the lateral edges and sometimes the callosities, scutellum except the outer angles of the basal lobe, and the elytra except the costal margin, more or less deeply infuscated, sometimes becoming a blackish-fuscous; disk of the front at times infuscated, the clypeus usually black and polished. Cuneus usually pale or decolorated. Membrane deeply infuscated with pale nervures and a small black ray beyond the tip of the areoles. Antennæ black with pale incisures, often with a broad pale annulus on the second joint. Legs pale, the femora more or less dotted with fuscous; the knees, tips of the tibiæ and the tarsi black. Above clothed with long gray pile giving the insect a hoary aspect.

Described from numerous examples taken about the lower end of Fallen Leaf Lake, California, in July, 1915.