

2. Pronotum with a distinct apical collar, or with a flattened anterior margin simulating a collar 3
 — Pronotum without a collar 9
3. Pronotal collar convex, separated from anterior disk by a distinct incised line 4
 — Pronotal collar flat, without an incised line behind it, or wanting; body elongated, often linear 8
4. Head viewed from above short, vertical, produced below the eye for nearly twice the length of the eye; antennae long and slender, much longer than the entire body, inserted the length of the clypeus above its base; basal joint of the tarsi as long as the following two together; arolia wanting; vertex deeply sulcate; pronotal collar very slender **Cylapini**
- Head not greatly produced below the eye; antennae rarely longer than the entire body, inserted about on the line of the base of the clypeus; basal joint of the tarsi shorter than the following two taken together 5
5. Rostrum long, passing the middle of the venter; head produced, horizontal or nearly so; tibiae smooth, or with minute pubescence only; tarsi slender, basal point but little shorter than the following two together; arolia wanting **Fulvini**
- Rostrum shorter, scarcely surpassing the hind coxae; tibiae armed with bristles or clothed with longer hair, rarely smooth 6
6. Pronotal collar broad, convex, about as broad as the callosities; tibiae thickly clothed with soft hairs but without rows of stouter bristles; body opaque, black, marked with red or fulvous **Horistini**
- Pronotal collar narrow, convex, often linear, rarely broad; then flat with the body elongated, tibiae smooth and pronotum broadest before the middle 7
7. Body elongated, often linear; base of scutellum usually exposed; tibiae smooth or nearly so; arolia minute and united with base of the claws, or as long as the claws and lying close to them **Dicyphini**
- Body rarely elongated, with the base of scutellum exposed and tibiae smooth or nearly so, arolia in this case free and divergent at apex **Capsini**
8. Tarsi long, first joint longer than the following two together; pronotal collar a mere flattening of the anterior margin, or sometimes wanting; vertex often sulcate; arolia large, free, often clavate **Mirini**
- Tarsi shorter, first joint not longer than the third, usually shorter, pronotal collar often wanting; arolia united with the claws or wanting **Hallodapini**
9. Arolia free, parallel, or converging toward their tips; wing-cell without a hamus **Orthotylini**
- Arolia wanting, or parallel with and usually united to the claws at base, wing-cell normally with a hamus **Phylini**