

13. Hind femora not at all saltatorial, not wider than the eye viewed from the side, pale, dotted in longitudinal series; tibial bristles black and inserted in black points; first joint of rostrum scarcely attaining base of head14
 — Hind femora saltatorial, much thickened or flattened, distinctly wider than lateral width of the eye15
14. Head broad, its width three-fourths that of hind margin of pronotum; viewed from the side produced below the eye for a distance as great as length of eye12. **Bolteria** Uhl.
 — Head narrower, hardly more than half as broad as basal width of the pronotum; viewed from the side shorter, produced below the eye for hardly more than half length of the eye; body densely clothed with white deciduous hairs11. **Oligotyus** n. gen.
15. Femora black, tarsi annulated with white; body small, black
20. **Strophopoda** n. gen.
 — Femora pale, dotted with fuscous or black points16
16. Base of clypeus on a line with the insertion of antennae
16. **Atomoscelis** Reut.
 — Base of clypeus above line connecting base of antennae
17. **Europiella** Reut.
17. Third joint of hind tarsi as long as first and second together; arolia short, linear; attached to base of claws.....18. **Tuponia** Reut.
 — Third joint of hind tarsi subequal to second, or shorter18
18. Head broad, about one-fifth narrower than hind margin of pronotum; body black, hind femora black19. **Chlamydatus** Curt.
 — Head narrower; body, including legs, pale19
19. Head viewed from the side forming a squarish projection below the eyes; legs and antennae pale, immaculate; tibial spines black; body pale, clothed with short, thick black hair...23. **Maurodactylus** Reut.
 — Head forming a very short angular projection below the eyes.....20
20. Legs pale, femora immaculate; second antennal joint annulate with black22. **Cylloceps** Uhl
 — Legs pale, hind femora dotted with black; first antennal joint annulate with black21. **Campylomma** Fieb.
21. Second antennal joint terete, but little thickened at apex, longer than third and fourth together24. **Rhinacloa** Reut.
 — Second antennal joint of the male broadly flattened22
22. Body black, elytra pale brown, bifasciate with white
14. **Leucopoecila** Reut.
 — Body entirely blue-black, or only coxae pale.....13. **Atractotomus** Fieb.