

2. **Monalonion versicolor**, n. sp. (Tab. XXIV. fig. 12.)

Head and antennæ shining black, the antennæ pilose, the head obscurely marked with castaneous near base; pronotum ochraceous, the whole of the anterior area and the lateral margins narrowly black, the ochraceous area containing a more or less arcuate castaneous fascia; scutellum ochraceous, its base opaquely greyish; corium pale reddish castaneous, the costal margins and a triangular spot in front of inner apical angles luteous, inner claval margins somewhat darker; cuneus reddish, the apex infuscated; membrane very pale fuscous and semihyaline, the cellular veins reddish; anterior legs black, intermediate and posterior legs ochraceous, the femora streaked with fuscous above, and the apices of the tarsi black. Body beneath *apparently* (carded specimen) with the head and sternum blackish, the abdomen ochraceous.

Long. 8 millim.

*Hab.* GUATEMALA, Las Mercedes (*Champion*).

3. **Monalonion hilaratum**, n. sp. (Tab. XXIV. fig. 13.)

Head black, with the base obscurely ochraceous; antennæ black, pilose, first joint with base and apex dull ochraceous, fourth joint and apex of third more or less fuscous; pronotum ochraceous, somewhat darker near lateral angles; scutellum reddish ochraceous; corium pale fuscous, the costal margins narrowly, at base broadly, ochraceous; cuneus bright reddish; membrane pale fuscous, the cellular veins reddish; legs ochraceous; tarsi and bases of posterior femora dark fuscous. Body beneath *apparently* (carded specimen) more or less reddish ochraceous.

Long. 9 millim.

*Hab.* GUATEMALA, San Gerónimo (*Champion*).

b. *Posterior tibiæ curved.*

4. **Monalonion atratum**, n. sp. (Tab. XXIV. fig. 14.)

Head, antennæ, and pronotum dark shining black, the antennæ strongly pilose; eyes luteous; scutellum and corium dark shining fuscous, the first with its apex somewhat paler, the second with the costal margin (particularly broad at base) luteous; membrane pale fuscous; legs black and strongly pilose, the anterior and intermediate tibiæ very broadly annulated with luteous. Body beneath *apparently* (carded specimen) shining black; abdomen above dull reddish.

Long. 10 millim.

*Hab.* PANAMA, Bugaba (*Champion*).

c. *Pronotum excavated at base.*

5. **Monalonion dissimulatum**, n. sp. (Tab. XXIV. fig. 15.)

Body above ochraceous; head, antennæ, anterior constricted area of pronotum, apex of scutellum, and a broad transverse fascia at apex of corium black; base of first joint of antennæ ochraceous; membrane pale ochraceous and semihyaline, the margins narrowly, and the apex broadly, fuscous; legs ochraceous; the tarsi, bases of posterior femora, and posterior tibiæ (excluding base) black. Body beneath with the sternum ochraceous; the head, anterior margin of prosternum, acetabula, and coxæ black; abdomen mutilated.

Pronotum ovately excavated at base.

Long. 11 millim.

*Hab.* GUATEMALA, San Isidro (*Champion*).