

20. **Resthenia montivaga**, n. sp.

Body above, legs, antennæ, and rostrum black, the basal half of the posterior femora dull ochraceous, and the third and fourth joints of the antennæ pale fuscous; bases of the first joint ochraceous. Sternum and abdomen beneath reddish ochraceous; disk of sternum and lateral margins of prosternum blackish; coxæ reddish ochraceous, but more or less suffused with blackish.

Long. 7 millim.

Hab. PANAMA, Volcan de Chiriqui 2500 to 4000 feet (*Champion*).

This species is allied to *R. atrata* (previously described at page 254) in superficial coloration; but differs in having a much more elongate and narrow body, the basal joint of the antennæ shorter, &c.

21. **Resthenia sudatus**, n. sp. (Tab. XXII. fig. 17.)

♀. Body above red; apex of head, antennæ, clavus (excluding basal third), apical margins of corium, cuneus, legs, central disk of mesonotum, and oviposital sheath to abdomen dark fuscous or black; basal vertex of head, a sinuated discal line near anterior margin of pronotum, margins of lateral pronotal angles, a sub-claval linear streak to corium, membrane and rostrum fuscous, the last with the apical margins paler; coxæ red, more or less suffused with fuscous.

The body is broad and subdepressed, the tibiæ somewhat sparingly setose, the antennæ obscurely pilose. Long. 8 millim.

Hab. MEXICO (*Koppe, Mus. Berol.*).

b. *Rostrum reaching or passing the posterior coxæ.*

c. *Basal joint of the antennæ appreciably longer than the head.*

22. **Resthenia bicolor**, n. sp.

Reddish ochraceous; eyes, basal half of pronotum (excluding lateral margins), scutellum, corium, membrane, apex of head, a spot at abdominal apex, tibiæ and tarsi fuscous or black; antennæ fuscous, the first joint luteous, with its base and apex infuscated; head with a central longitudinal fuscous line on vertex; rostrum obscure reddish or pale fuscous, the basal joint reddish ochraceous and the apex pithy.

Antennæ somewhat thickly and strongly pilose; tibiæ strongly and thickly setose. Long. 8½ to 9 millim.

Hab. GUATEMALA, Zapote (*Champion*).

cc. *Basal joint of the antennæ about equal in length to that of the head, or very little longer or shorter.*

23. **Resthenia bivittis**. (Tab. XXIV. fig. 21.)

Resthenia bivittis, Stål, Stett. ent. Zeit. xxiii. p. 318. 239¹.

Hab. MEXICO¹.

The type in the collection of Dr. Signoret is here figured.

24. **Resthenia vitticeps**. (Tab. XXIV. fig. 22.)

Resthenia vitticeps, Stål, Stett. ent. Zeit. xxiii. p. 318. 242¹.

Hab. MEXICO¹.

I am again indebted to Dr. Signoret for the loan of the type, which is here figured.