

a. *Body subelongate. Pronotum not prominently gibbous.*

1. ***Eccritotarsus generosus***. (Tab. XXII. fig. 24.)

Eccritotarsus generosus, Stål, Stett. ent. Zeit. xxiii. p. 323. 254¹.

Eccritotarsus eucosmus, Stål, Stett. ent. Zeit. xxiii. p. 323. 255².

Hab. MEXICO^{1 2} (*coll. Sign.*); GUATEMALA, Zapote, San Isidro (*Champion*),

A typical specimen of *E. generosus* in the Stockholm Museum is here figured, with which I have compared typical specimens of *E. eucosmus* in the collection of Dr. Signoret, and have come to the conclusion that differential characters do not exist, sufficient at least to denote more than one species.

2. ***Eccritotarsus nigripes***, n. sp.

Dark bluish black; head (both above and beneath) and pronotum sanguineous; the eyes vary in colour from sanguineous to greyish ochraceous; the body above is distinctly and finely punctate.

Var. a. Basal joint of the antennæ and basal margin of the scutellum sanguineous.

Long. 4½ millim.

Hab. MEXICO, Ciudad (*Forrer*); GUATEMALA, Cubilguitz (*Champion*).

3. ***Eccritotarsus genetivus***, n. sp. (Tab. XXII. fig. 25.)

Head and pronotum reddish ochraceous; scutellum, corium, and body beneath piceous; membrane fuscous; coxæ and legs luteous, apices of the posterior femora and apices of the tarsi fuscous; rostrum luteous, its apex piceous; antennæ with the basal joint ochraceous; second joint piceous, remaining joints mutilated.

Long. 4½ millim.

Hab. MEXICO (*Deppe, Mus. Berol.*).

4. ***Eccritotarsus splendens***, n. sp. (Tab. XXVIII. fig. 1.)

Dark shining bluish; head, pronotum, sternum, and legs luteous; antennæ piceous, base of the first joint luteous; rostrum luteous, with its apex piceous; eyes blackish; legs with an upper linear streak to anterior femora, a subapical broad annulation to intermediate and posterior femora, basal and central annulations to tibiæ, and base and apex of tarsi fuscous; membrane smoky subhyaline, its basal cellular area piceous.

Long. 4½ millim.

Hab. PANAMA, Bugaba (*Champion*).

In this species the eyes are large and considerably recurved; the antennal joints are uniform, and not distinctly incrassated.

5. ***Eccritotarsus vestitus***, n. sp. (Tab. XXVIII. fig. 2.)

Ochraceous; antennæ, membrane, tibiæ, and tarsi piceous; scutellum, clavus (excluding base), and inner margins of corium bluish black; apical margin of the membrane smoky hyaline. The ochraceous colouring of the corium is more rufous than that of the head and pronotum.

Long. 4½ millim.